

Pattern of Psychiatric illness among Tuberculosis Patients

*AT Islam¹, M Hussain², MA Siddique³, SM Badruddoza⁴

¹Dr. Ahmed Tanjumul Islam, Medical Officer, Department of Medicine, Rajshahi Medical College Hospital,

²Dr. Mamun Hussain. Associate professor, Department of Psychiatry, Rajshahi Medical College Hospital

³Dr. Md. Abdullah Siddique, Professor, Department of Microbiology, Rajshahi Medical College Hospital

⁴Dr. Shah Md. Badruddoza, Professor, Department of Pathology, Rajshahi Medical College Hospital

*Corresponding Author

Date of submission: 01 April 2015

Date of acceptance: 05 May 2015

ABSTRACT

Background: The incidence of tuberculosis is very high in Bangladesh. There is a high prevalence of psychiatric illness among tuberculosis patients. But primary care physicians and specialists do not screen this association. The aims of this study were to evaluate the incidence and pattern of psychiatric illness in tuberculosis patients.

Methodology: The study population included those patients who were coming for treatment in DOTs corner of Rajshahi Medical College Hospital between July 2014 - February 2015 (Eight months). Each patient underwent a detailed psychiatric evaluation by a consultant psychiatrist once they were medically stable. Details included socio-demographic data, psychiatric diagnosis and treatment outcome. All data was tabulated and analyzed using SPSS-16.

Result: Among the total 350 tuberculosis patients, total 108 patients (31%) were diagnosed with psychiatric problems. Most of them are male (71%), less than 40 years of age (70%) with no previous psychiatric illness (98%). Depression (n=43, 40%) and Anxiety (n=33, 31%) were the commonest psychiatric illness diagnosed. Fear of disease outcome was the commonest precipitating factor found (66%). After the psychiatric treatment, 94% improved clinically.

Conclusion: Detecting the level of psychiatric illness among tuberculosis patients at early stage will improve continuation and adherence to treatment. A referral system to psychiatrists by physicians needs to develop to screen the mental disorder symptoms to treat these co morbidities.

Key Words: Tuberculosis, Psychiatric illness, Bangladesh

Introduction

Tuberculosis (TB) is a chronic infectious multi systemic disease caused by mycobacterium tuberculosis and is one of the leading causes of mortality worldwide¹. The World Health Organization (WHO) has estimated that 2 billion people, almost a third of the world's population have latent tuberculosis². Every year about eight million people develop this disease, and some three million die of it, over 95% of these from developing countries. In 2005 the highest rates per capital were from Africa (28% of all TB cases), and half of all new cases were from five Asian countries, namely Bangladesh, China, India, Indonesia and Pakistan^{3, 4}. Tuberculosis mortality is an important indicator of the success of TB control. A higher rate of mortality and morbidity was seen among patients with baseline

psychiatric, because they defaulted from treatment⁵. There is a high prevalence of psychiatric illness in TB patients, but primary care physicians and specialists do not screen this association although anxiety and depression occur frequently in persons with these cases.

It has also been shown in most of the National and International studies that most of these patients have a history of mental illness; the commonest diagnosis being depression which is usually followed by personality disorder alone or co morbid with other psychiatric illnesses⁶.

Keeping these facts and figures in mind, we intended to conduct a study aiming to analyze the patterns and determinants of psychiatric illnesses in the patients diagnosed as tuberculosis in a tertiary care hospital.

Materials and Methods

This cross-sectional descriptive study based on the interview of the patients diagnosed as tuberculosis comes for treatment in DOTs (Directly Observed Therapy) corner. The aim of this study was to evaluate the pattern of psychiatric illness in tuberculosis patients at early stage and subsequently how it can improve continuation and adherence to treatment.

The study population included all those patients who were diagnosed as tuberculosis and being managed for and brought for psychiatric evaluation during the period of eight months (July 2014- February 2015). Each patient underwent a detailed psychiatric evaluation by a consultant psychiatrist. Psychiatric diagnoses were considered as per ICD-10 criteria and patients were managed with pharmacological / non-pharmacological measures. The factors related to suicide (intentionality and lethality) were assessed during the interview and mental status examination of the patients. Besides the questionnaire, level of depression was rated by BDI, and anxiety was rated by STAI.

The Beck Depression Inventory (BDI) is questionnaire for measure the intensity, severity, and depth of depression. It is composed of 21 questions. Clinically significant level of depression was defined by score of BDI: 0-4 Normal, 5-7 mild, 8-15 moderate, >16 severe Depression.

STAI, or State-Trait Anxiety Inventory, is an instrument that quantifies adult anxiety. It is questionnaire used to simplify the separation between state anxiety and trait anxiety, feelings of anxiety and depression. We used S-Anxiety scale STAI ST and the T-Anxiety scale STAI TR, each having 20 items. These tests are answered on the basis of a 1-4 scale, with the focused areas including: worry, tension, apprehension, and nervousness. Value 40-60 interpreted like moderate and more than 60 sever symptoms.

Data was tabulated and statistical analysis was performed using software SPSS-16.

Results

Among the 350 patients diagnosed as tuberculosis, 108 patients (31%) were found with different patterns of psychiatric illness. Four types of tuberculosis diagnosed with psychiatric illness in our study. Of them 80 new cases found (74%). 11, 9 and 3 cases were found as relapse, treatment failure and multidrug resistant tuberculosis (MDR) respectively. (Figure 1)




Figure-1: Tuberculosis types with psychiatric illness.

On studying socio-demographic factors among tuberculosis patients we found 70% patients were aged less than 40 years. 71% affected individual were male. 49% were married. 51% were from urban area. 66% were employed. At the time of interview smoker (43%) and alcoholic (12%) patients also found. 32% had co morbid illness predominantly with hypertension (14%) and diabetes (13%). Only 2% had previous psychiatric illness. (Table 1)

Table 1: Distribution of Socio-demographic factors in Tuberculosis patients:

Age Group	Number (n)	Percentage %
<40 years	76	70%
>40 years	32	30%
Sex group		
Male	77	71%
Female	31	29%
Marital status		
Single	49	46%
Married	53	49%
Widowed	06	5%
Economic Status		
Employed	71	66%
Unemployed	37	34%
Residence		
Rural	53	49%
Urban	55	51%
Substance use:		
Smoker	46	43%
Alcoholic	13	12%
Nothing	43	40%
Others	6	5%
Co morbid disorders		
None	73	68%
Hypertension	15	14%
Diabetes	14	13%
Ischemic Heart disease	5	4%
HIV	1	1%
Past psychiatric illness		
Yes	3	2%
No	105	98%

In our study 108 patients were diagnosed with psychiatric illness among 350 tuberculosis patients making the prevalence of psychiatric illness 31%. (Figure 2)




Figure 2: Percentage of Psychiatric illness in Tuberculosis patients.

Various types of Psychiatric illness were found in patients who were diagnosed as tuberculosis. Of the total 108 cases seven types of psychiatric illness were found. Of them Depressive illness was the commonest (40%, n=43). Others are anxiety (n=33), somatoform (n=11), psychotic spectrum disorder (n=9), personality disorder (n=7) and substance use disorder (n=3). 2 cases remained undiagnosed. (Figure 3)




Figure 3: Number and Types of Psychiatric illness among tuberculosis patients.

We also found that Depression (40%, n= 43) and Anxiety (31%, n=33) were the most common psychiatric findings (71%) among tuberculosis patients. Only 29% diagnosed as other psychiatric illness. 2 cases remain undiagnosed. (Figure 4)




Figure 4: Common Psychiatric findings in TB patients.

There were multiple psychosocial precipitating factors among tuberculosis patients.

Fear of disease outcome (66%, n=71) and length of treatment (44%, n=48) were the most common precipitating factors. Current Health status (29%), social issues (27%), family issues (18%), financial issues (17%), side effects of drugs (8%) also identified as precipitating factors. (Figure 5)




Figure 5: Psychosocial precipitating factors among tuberculosis patients

*Each person allowed to answer multiple factor.

We also studied patient's mental perspective about illness. 76% patients knew about their diagnosis. 17% thinks the disease as malignancy. 5% did not know the diagnosis. 56% knows correctly why the disease occurs. 56% were worried about their families and their children's future (40%). 66% expects to cure the disease fully. (Table 2)

Table 2: Patients perspective about their illness.

*Allowed to give multiple answers.

Belief about illness	Number	Percentage%
<i>Name of main problem</i>		
TB	81	76 %
AIDS	3	2 %
Carcinoma	18	17 %
Don't know	6	5 %
<i>Causation</i>		
Germs	61	57 %
Poor diet	10	9 %
Smoking	16	15 %
Alcohol	9	8 %
Don't know	12	11 %
<i>Fear about illness*</i>		
Death	41	38 %
Disease will worsen	30	28 %
Will infect family	61	57 %
Worry children's future	4 ₃	40 %
Others	10	9 %
<i>Treatment expectations</i>		
Will be fully cured	71	67 %
Not fully cured	2 ⁴	22 %
Lead to disability	5	4 %
Don't know	8	7 %

Our study also reveals distribution of treatment related factors. Pharmacologic treatment were given with Benzodiazepines (36%, n=36%), antidepressants (29%, n=29%), antipsychotics (8%, n=9). Non pharmacologic treatment (counseling) were given to 21 %. Outcome was excellent with 94% improved or recovered clinically. (Table 3)

Table 3: Distribution of Treatment related factors:

Treatment given (n=108)	Number	Percentage %
Benzodiazepines	39	37%
Antidepressants	31	29 %
Antipsychotics	9	8 %
Combined	6	5 %
Counseling	23	21 %
<i>Outcome</i>		
Improved/ Recovered	101	93%
Treatment failure	5	3%
Not returned	4	3 %
Expired	1	1 %

Discussion

Tuberculosis is a common disease in developing countries causing high mortality and morbidity¹. Due to its long treatment process and variable outcomes psychiatric illness is common in tuberculosis patients^{2,3}. Psychiatric complications (anxiety, depression, psychosis) can greatly impact patient quality of life of patients with tuberculosis specially MDR- TB therapy^{6,8}. In our study we have shown that Depression and anxiety are very high in patients with tuberculosis (Depression 42 %, Anxiety 29%). Depression was positively correlated with anxiety ($p=0.001$) for patient with tuberculosis.

Baba and his colleagues showed 7% prevalence of depression in Nigeria⁵. A study conducted in India showed that the prevalence of depression and anxiety was 19% for recently diagnosed patients with TB, 22% for defaulted TB patients and 26% for patients with Multidrug resistant¹⁰. A higher prevalence of psychiatric disorders was found in tuberculosis group (31%) compared with 5% in healthy control group in Nigerian study⁵. The difference of prevalence rates can be explained possibly by sensitivity of screening instruments used, other psychological factors associated with hospital admission, educational level, other co morbidities and complications of tuberculosis. In United Kingdom a higher rate of depression and anxiety were in noncompliant TB patients and treating psychological disorders may substantially improve treatment adherence⁴. The tuberculosis patients with an unfavorable social status with stressful economic factors (financial worry, unemployment, homeless, poverty) and stressful personal factors (divorce or alone, multiple treatment failures or abandon) also carries high risk for psychiatric illness⁷.

Raised depressive prevalence (46%) and anxiety scores (47%) were associated with an increase in the number of symptoms reported, more serious perceived consequences and less control over the illness⁶. Persistent cough may be an additional control of depressive patient³.

The patients' perspectives of their illness showed that a significant proportion of the subjects did not believe in the infectious nature of the disease and feared death. These finding are similar to the studies reported from other developing nations¹⁰, despite the health education. A detailed elicitation of the perspectives held by each patient, tackling these

beliefs at the individual's level rather than health education in groups, and in a culturally sensitive way taking into account the local perceptions of the illness is indicated. Different studies also showed psychiatric illness as an important factor for poor medication compliance in tuberculosis^{10,11}.

Conclusion

Evaluation and management of mental disorders from Tuberculosis Patients (significantly higher compared to general population) may increase treatment compliance and reduce relapse. This can ameliorate the prognosis and quality of life for patients with this chronic disease. Primary care doctors and consultants need to develop systematic strategies to screen the mental disorders symptoms in tuberculosis patients and must ask aide from psychology or psychiatric doctor for treat these co morbidities.

Conflict of interest: we have no conflict of interest

References

1. Grange JM, zumla A. Paradox of the global emergency of tuberculosis. *Lancet* 2009. 353:996
2. Kochi A. Tuberculosis control - Is DOTS the health breakthrough of the 1990s? *World Health Forum* 2001; 18:225
3. Moulding TS. Medication monitors to treat tuberculosis, A supplement to directly observed therapy, *Am J Respir Crit Cure Mecl* 2001; 159:989
4. Falab-Stubi CL, Zellweger JP. Sauty A, Uldry C, Iorillo D, Burner M. Electronic monitoring of adherence to treatment in the preventive chemotherapy of tuberculosis *Int J Tuberc Luni > Di.* 1998; 2:525
5. Baba A Issa, Abdullah D Yussuf, Suleiman I Kuranga: Depression comorbidity among patients with tuberculosis in a university teaching hospital outpatient clinic in Nigeria. *Mental Health Family Medicine*, 2009 September; 6(3): 133-138
6. Vega P, Sweetland A, Acha J, Castillo H: Psychiatric issues in the management of patients with multidrugresistant tuberculosis. *Int. J. Tuberc. Lung Dis.* 2004 jun. 8 (6):749-759.
7. Thomas H. Holmes, Norman G. Hawkins, Charles E. Bowerman, Edmund R. Clarke , Joy R. Joffe : Psychosocial and Psychophysiologic Studies of Tuberculosis. *Psychosomatic Medicine* 19:134-143 (1957)
8. L. Lewis GH, Pelosi A, Araya R. Dunn G. Measuring psychiatric disorder in the community: a standardised assessment for lay interviewers. *Psychol Mecl* 2009; 22:465
9. Lloyd KR, Jacob KS, Patel V. St. Louis L, Bhugra D, Mann A. The development of the Short Explanatory Model Interview (SEMI) and its use among primary-care attenders with common mental disorders. *Psychol Med* 2001; 28:1231
10. Cornolet TM, Rakotomalala R. Rajaonarioa H. Factors determining compliance with tuberculosis treatment in an urban environment, Tamatave, Madagascar. *Int J Tuherc Luna Dis* 2002; 2:891
11. Menzies R, Rocher I, Vissandjee B. Factors associated with compliance in treatment of tuberculosis. *Tubercle & lung Dis* 1993; 74:32