Case Reports

Congenital Hypotrichosis Simplex: A Case Report

MARIA KIBTIAR¹, ROMANA AKTHER¹, MOHAMMED MAHBUBUL ISLAM², SHAHANA A RAHMAN³

Abstract:

Congenital hypotrichosis simplex of scalp is a rare disorder which only affect the scalp hair. Rest of the hair bearing areas are spared. In this disorder the scalp hair are absent or sparse. We describe a 5 month old boy presented with no growth of scalp hair since birth in this case report. After taking detailed history, a thorough physical examination was done. Other causes of alopecia were ruled out and finally the child was diagnosed as a case of Congenital Hypotrichosis Simplex (CHS).

Introduction:

Congenital Hypotrichosis Simplex of scalp is a genetic disorder characterized by sparse or absent scalp hair without structural defect of the hair and absence of other systemic abnormalities¹. Prevalence of the disease is <1/100000 population and male and female are equally affected². Both autosomal dominant and recessive modes of transmission have been reported in this disorder². In some patients the diagnosis of sporadic hypotrichosis simplex should be considered after rulling out all other possible causes of congenital hypotrichosis³. There is no abnormalities in other ectodermal derrivatives, so skin, nail and teeth remain normal in these patients⁴. There is no specific treatment of this disease . Due to variable expression of the disease proper counselling is of great importance⁵.

Case report:

A 5 month old boy, 2nd issue of non consanguineous parents was admitted at the Department of Paediatrics, Bangabandhu Sheikh Mujib Medical university (BSMMU) with the complaints of no hair growth since birth along with some itchy erythematous skin rashes over the scalp. His perspiration was normal and his eyebrow and nail growth were also normal. Other family members were healthy with no evidence of such type of illness.

- 1. Resident phase-B, Department of Paediatrics, Bangabandhu Sheikh Mujib Medical University, Dhaka.
- 2. Assistant Professor, Department of Paediatrics, Bangabandhu Sheikh Mujib Medical University, Dhaka.
- 3. Professor of Paediatrics & Pro Vice Chancellor (Academic), Bangabandhu Sheikh Mujib Medical University, Dhaka.

Correspondence: Dr. Maria Kibtiar, Resident phase-B, Department of Paediatrics, Bangabandhu Sheikh Mujib Medical University, Dhaka. Email: mariabsmmu@yahoo.com, Mob: 01710327674.

Received: 24-06-2018

Accepted: 16-03-2019



Fig.-1: No scalp hair

On examination there was no scalp hair (Figure 1) but his eyebrows, eyelashes and nails were normal (Figure 2 and 3). There were some maculopaular rashes present over the scalp. The child was developmentally age appropriate. Examination of other system revealed normal findings. Provisionally he was diagnosed as a case of Congenital Hypotrichosis Symplex (CHS).

BANGLADESH J CHILD HEALTH 2019; VOL 43 (1) : 60



Fig.-2: Normal nails in feet

Work out and investigations were done to rule out congenital hypothyroidism and diagnose the case which included normal T4 and TSH level. Neurometabolic diseases were also excluded as S.ammonia ,S.lactate, urinary ketone body and urinary PH were normal. Scalp biopsy was done which revealed perifollicular fibrosis and mild perifollicular infiltration of chronic inflammatory cells which were consistant with Congenital hypotrichosis simplex (CHS). Finally the baby was diagnosed as Congenital hypotrichosis simplex(CHS).

Discussion:

Congenital hypotrichosis simplex of scalp is a rare autosomal dominant genotrichosis characterized by a hair defect confined to scalp in the absence of other ectodermal and systemic abnormalities⁴. In this disorder other ectodermal derivatives such as eyebrows, eyelashes, nails and sweat glands remain normal⁴. Rafique et al. in their study showed that the genetic defect was located to chromosome 18g21.1⁶. Shimomura Y et al. identified a mutation in the adenomatosis polyposis down-regulated 1(APCDD1) gene in CHS⁷. If there is presence of strong family history the diagnosis is straight forward but in isolated cases like our one the diagnosis of sporadic hypotrichosis simplex of scalp can be done only by ruling out other possible causes of congenital and hereditary hypotrichosis.

Ectodermal dysplasia which is one of the important cause of hypotrichosis was excluded in our patient



Fig.-3: Normal nails in hands

as the eyelashes, nail and sweat glands were unaffected.

We also excluded congenital hypothyroidism and metabolic disorders where there may be hypotrichosis. Another rare congenital disorder is Marie Unna hypotrichosis where there is absence or scarcity of hair, eyebrows and eyelashes at birth, coarse and wiry hair during childhood and progressive hair loss begins around puberty⁸. But in our patient the eye brows and eyelashes were normal. CHS also need to be differentiated from aplasia cutis congenita, alopecia triangularis and congenital atricia⁹.

In aplasia cutis congenita baby is born with absent patch of skin resembling to open wound¹⁰. In our patient the skin was normal. In patient with alopecia triangularis a well defined triangular patch of scalp is affected¹¹. In our patient the whole scalp was affected. The congenital atrichia baby usually born with normal scalp hair but in early childhood hair is lost and never regrow again⁹ which was also not found in our patient as he had no hair growth since birth.

We also did scalp biopsy in our patient which showed perifollicular fibrosis with perifollocular infiltration of chronic inflammatory cells consistant with CHS. In a previous case report of hereditary hypotrichosis, the scalp biopsy showed marked decrease number of hair follicles accompanied by mild chronic predominantly lymphocyte perifollicular infiltration¹. Congenital Hypotrichosis Simplex: A Case Report

There is no effective treatment of CHS. Due to phenotypic variation patient need counselling⁵. We also counselled the parents about the disease during the course of admission and discharge. Besides that we prescribed an emollient for his rough skin over the scalp along with antihistamin syrup for itchy skin lesion.

Conclusion:

Congenital hypotrichosis simplex , a rare form of hypotrichosis has been largely regarded as a genetic disorder usually confined to the scalp. The disease can be diagnosed by excluding other possible causes of hypotrichosis. There is no specific treatment , counselling is the mainstay of management.As because it is a rare disease a good knowledge is required for early identification of the cases.

References:

- Vazquez MR, Rodiguez RR, topia AG, Diez L. Hereditary Hypotrichosis Simplex of the scalp. Pediatr Dematol 2002; 19: 148-150.
- 2. Orphanet Epidemiological Data: Congenital Hypotrichosis Simplex.2010. https:// www.orpha.net
- 3. Cambiaghi S, Barbareschi M. A sporadic case of Congenital Hypotrichosis Simplex of the

scalp: difficulties in diagnosis and classification. Pediatr Dermatol 1999; 16: 301-304.

- 4. Farshi HM, Ayatollahi A, Hejazi S.Heretidary hypotrichosis simplex. Indian J Dermatol 2014; 59:634.
- 5. Ronda S. Farah, Heather M, Holahan, Molly S, Moye et al. Hereditary Hypotrichosis Simplex of the Scalp. Cutis 2017; 100: 12-14.
- 6. Rafique MA, Ansar M, Jamal SM, Malik S, Sohail M, Haque MF et al. A locus for hereditary hypotrichosis localized to human chromosome 18q21.1. European Journal of Human Genetics 2003;11: 623-628.
- Shimomura Y, agalliu D, Vonica A, Luria V, wajid M, Baunner A. APCDD1 is a novel Wnt inhibitor mutated in hereditary hypotrichosis simplex. Nature 2010; 464 :1043-1047.
- 8. Shahana M, Hiremagalore R. Marie-Unna hereditary hypotrichosis.International Journal of Hypotrichosi 2014 ;6 :182-184.
- Naqash MM, Salati SA. Congenital hypotrichosis simplex. J Pioneer Med Sci 2015; 5: 190-191
- 10. Salati SA. Aplasia Cutis Congenita of Scalp. J surg park 2009; 14 :187-188.
- 11. Yin Li VC, Yesudian PD. Congenital triangular Alopecia; Int J Trichology 2015; 7 : 48-53.