

Minimal-invasive surgery for treatment of colo-rectal angiodysplasia in a child

A. Moreddu Eric¹
 Fabre Alexandre²
 Boubnova Julia¹
 Lamy Frederic³
 Guys Jean-Michel¹
 de Lagausie Pascal^{1*}

¹Department of Paediatric Surgery, Hospital La Timone, Marseilles, France

²Department of Paediatric Gastroenterology, Hospital La Timone, Marseilles, France

³Department of Anaesthesia/Intensive care, Hospital La Timone, Marseilles, France

*Correspondence to:

Pr de Lagausie Pascal, MD PhD
 Department of Paediatric Surgery
 Hospital La Timone, Marseilles
 264 rue Saint Pierre, 13385 Marseilles
 France
 E-mail: pascal.delagausie@ap-hm.fr
 Tel: +33(0) 491 386 679
 Fax: +33(0) 491 385 746

Summary

Introduction: Rectal angiodysplasia is a rare lesion of bowel wall in pediatric population. **Materials and Methods:** We report the first laparoscopic management of a case of bifocal angiodysplasia in a child, involving the rectum and the left colonic flexure. The preoperative investigations showed only rectal lesions. A classic Swenson's pull-through by laparoscopy was performed, with discovery of second lesion necessitating a resection of the left colonic flexure. **Result:** At day 11, the child presented an occlusive syndrome and a partial disunion of the coloanal anastomosis, requiring a discharge transverse colostomy. Two months after closure of the colostomy, the child regained 1 kg, had a normal transit (3 to 4 defecations per day) and normal haemoglobin results.

Key words: Colo-rectal; angiodysplasia; minimal-invasive surgery; venous malformations; laparoscopy; endorectal pull-through.

INTRODUCTION

Rectal angiodysplasia is a rare lesion of bowel wall in pediatric population, frequently revealed by bleeding.¹⁻⁵ Its diagnosis is difficult and made by the combination of multiple techniques.³⁻⁵ The usual treatment of colonic and rectal angiodysplasia is an open surgery, with resection of the affected zone.^{2,3} In rectal angiodysplasia, the classic technique is a Swenson pull-through with dissection of the rectum by an open abdominal surgery, also used to cure Hirschsprung's disease.³

It has been demonstrated in previous studies⁶ concerning Hirschsprung's disease that laparoscopic pull-through technique reduces complications, soiling and bleeding compared to open surgery,⁷ with earlier oral intake and discharge,⁸ but no significant difference about continence.⁹

To our knowledge, minimal-invasive surgery has not been yet described to cure angiodysplasia. This is the report of a case of rectal and colonic angiodysplasia with laparoscopic management on a young 4-year-old girl.

MATERIALS AND METHODS

A 4-year-old female infant was presented to the emergency department for massive rectal bleeding, with severe anaemia (haemoglobin at 3.4 g/dL).

How to cite this article: Moreddu E, Fabre A, Boubnova J, Lamy F, Guys JM, De Lagausie P. Minimalinvasive surgery for treatment of colo-rectal angiodysplasia in a child. Bang J Endosurg 2013;1(1): 13–15.




Figure 1: Pelvic MRI, T2 sequence: venous dilatations in the rectal wall

The oeso-gastro-duodenal fiber endoscopy and abdominal ultrasounds were normal. The rectoscopy demonstrated an erythematous rectal mucosa,² ulcerations of the rectum and an important vascular hyperplasia. A total colonoscopy was performed and did not show the presence of any vascular dysplasia's

The pelvic magnetic resonance imaging (MRI) demonstrated capillary and venous dilatations in the perirectal fat and in the rectal wall, making the diagnosis of rectal angiodysplasia (Fig. 1).

We decided a rectal surgery by Swenson's pull-through technique by laparoscopy. Four ports (three 5 mm for instrumentation and one 10 mm for telescope) were used.

The exploration of the colon revealed a 20-cm long angiodysplastic zone above the anus and, surprisingly, the same aspect on the left colonic flexure that had not been diagnosed on the preoperative tests. The rectum and the left colonic flexure were released.

A proctectomy with Swenson's pull-through (endo-rectal pull-through) technique was performed: circumferential section 2 mm above the ano-pectineal line, with trans-rectal dissection allowing moving down the colon and making a colo-anal anastomosis. The whole dysplastic zone of the rectum was removed during the intervention (Fig. 2).

After positioning two digestive clamps, a laparoscopic resection-anastomosis of the left colonic flexure was performed on the second angiodysplastic zone: intra-abdominal anastomosis (two 5/0 running sutures). The per-operative aspect seemed correct, with no traction on the colon.




Figure 2: Per-operative aspect of the rectum and the colon, after dissection and before coloanal anastomosis (black arrow)

RESULTS

The first post-operative days were simple with stool movements at day 3, and feeding at day 4. Unfortunately, at day 11, the child presented an occlusive syndrome and a partial disunion of the coloanal anastomosis, requiring a discharge transverse colostomy.

The histopathology confirmed the diagnosis of angiodysplasia. On the left colonic flexure, a moderate aspect of these lesions was described.

One month later, after a normal colonic anterograde opacification and satisfying digital rectal examination, the colostomy was closed.

Two months after closure of the colostomy, the child regained 1 kg, had a normal transit (3 to 4 defecations per day) and normal haemoglobin results. No more bleeding episodes occurred. Follow-up is now more than two years. Continence and transit are normal, with a few medication (SMECTA®).

DISCUSSION

To our knowledge, we report the first case of minimal-invasive surgery for rectal angiodysplasia. Its originality comes from the discovery of bifocal lesions, on the rectum and the left colonic flexure.

Minimal-invasive surgery allows a comprehensive exploration of the bowel. In this case, the discovery of the second

angiodysplastic zone on the left colonic flexure was made by laparoscopy, while the preoperative colonoscopy was normal.

The rectal wall dissection was easy and non hemorrhagic (no per-operative transfusion). This step was facilitated by laparoscopic magnification and probably most respectful of pelvic nervous structures.

The histopathologic results of the lesions in this case are similar to the reported cases in literature: venous dilatations in the sub mucosa.¹⁻³ In this case, the colonoscopy did not reveal the moderate lesions on the left flexure. Even if it is difficult to predict the evolution and bleeding of these lesions, a systematic removal is indicated.

The laparoscopic surgery was easily performed, but the postoperative course was complicated. Traction on both anastomoses are probably responsible for these complications. In our mind, they could have been avoided by a two-time procedure: a first time with Swenson's technique, and a second time, after healing, on the left colonic flexure. With this two time procedure, we probably decrease the risk of leakage. A single time treatment for multifocal lesions is not recommended, and particularly in this situation, where the lesions remain close to each other.

A complete exploration of the colon is necessary since multiple localizations are possible and can be unseen by preoperative tests.⁴ Laparoscopy is an interesting way for this global bowel exploration.

CONCLUSION

Laparoscopy is a good approach for pull-through in the treatment of angiodysplasia. We recommend it, because

more than the advantages of a minimal invasive surgery, it allows a global sight of bowel lesions.

References

1. Ayadi A, Pousse H, Soua H, et al. A rare cause of rectal hemorrhage in children: rectal angiodysplasia. *Arch Pediatr*. 1994 May;1(5):493-6.
2. de la Salle R, Aubert D, Nachin P, Pageaut G, Gille P. Daily rectal bleeding caused by venous angiodysplasia of the rectum. *Chir Pediatr*. 1984;25(3):189-90.
3. de la Torre Mondragón L, Vargas Gómez MA, Mora Tiscarreño MA, Ramírez Mayans J. Angiodysplasia of the colon in children. *J Pediatr Surg*. 1995 Jan;30(1):72-5.
4. Ozer I, Birol Bostanci E, Ozogul YB, et al. Diffuse colorectal angiodysplasia misdiagnosed preoperatively as ulcerative colitis in a child: report of a case. *Surg Today*. 2008;38(8):731-3.
5. Soon-Ok Choi, Woo-Hyun Park, Jin-Bok Hwang, Sang Pyo Kim, Ae Suk Kim. Diffuse microscopic angiodysplasia and nodular lymphoid hyperplasia in an infant with obscure massive lower gastrointestinal bleeding: a diagnostic challenge. *J Pediatr Surg*. 2009 April;44(4):852-5.
6. de Lagausie P, Bruneau B, Besnard M, Jaby O, Aigrain Y. Definitive treatment of Hirschsprung's disease with a laparoscopic Duhamel pull-through procedure in childhood. *Surg Laparosc Endosc*. 1998 Feb;8(1):55-7.
7. Kubota A, Kawahara H, Okuyama H, Oue T, Tazuke Y, Okada A. Clinical outcome of laparoscopically assisted endorectal pull-through in Hirschsprung's disease: comparison of abdominal and perineal approaches. *J Pediatr Surg*. 2004 December;39(12):1835-7.
8. Curran TJ, Raffensperger JG. Laparoscopic Swenson pull-through: a comparison with the open procedure. *J Pediatr Surg*. 1996 August;31(8):1155-7.
9. Fujiwara N, Kaneyama K, Okazaki T, Geoffrey JL, Kato Y, Kobayashi H, Yamataka A. A comparative study of laparoscopy-assisted pull-through and open pull-through for Hirschsprung's disease with special reference to postoperative fecal continence. *J Pediatr Surg*. 2007 December;42(12):2071-4.