

# DIFFUSE OLIGOSYMPOMATIC CAROLI'S DISEASE: CASE REPORT

LEMFADLI Y<sup>1</sup>, BOUHRIT S<sup>1</sup>, AIT ERRAMI A<sup>1</sup>, OUBAHA S<sup>2</sup>, SAMLANI Z<sup>1</sup>, KRATI K<sup>1</sup>

## Abstract

*This article describes a case of Caroli's disease in a 53-year-old female patient who complained non-specific abdominal pain without cholestasis or cholangitis. Ultrasound and hepatic magnetic resonance imaging showed segmental saccular dilations connected to intrahepatic bile ducts without hepatic fibrosis. This clinical case shows the possibility of having oligosymptomatic forms in the diffuse forms of Caroli disease, therefore the interest to consider this diagnosis in case of non-specific abdominal signs and to request a hepatic ultrasound.*

**Keywords:** Caroli disease, Caroli syndrome, congenital bile duct dilatation.

Received: 16 April 2020

Accepted: 12 June 2020

DOI: <https://doi.org/10.3329/bjm.v31i2.48540>

## Introduction

Caroli's disease (CD) is a rare congenital disease, first described in 1958 by Jacques Caroli.<sup>1</sup> It is part of the spectrum of fibropolycystic liver pathologies. It is characterized by congenital non-obstructive dilation of the intrahepatic bile ducts. It can be localized or diffuse. Caroli syndrome is defined as its association with congenital hepatic fibrosis.<sup>2</sup> It is most often revealed by recurrent episodes of cholangitis.<sup>3</sup>

This article describes the case of CD in a 53-year-old female patient with cystic formations distributed throughout the hepatic parenchyma, fortuitously diagnosed in adulthood, during the etiological investigation of non-specific abdominal pain.

## Case report

A 53-year-old female patient was seen in consultation. She complained of abdominal pain, which had started a year ago. The intensity of the pain was mild to moderate, intermittent, recurrent, located in the upper right quadrant and relieved by taking first-level pain relievers. The patient also reported urinary burns, but no fever or chills. She did not report jaundice, pruritus, vomiting, or weight loss.

The patient had no history of personal or family hepatopathy. She was taking no treatment and no toxic habits.

The clinical examination, in particular the abdominal, respiratory and cardiovascular examination, did not show any abnormalities. There was no jaundice or fever. The abdominal examination did not show hepatomegaly or signs of hepatocellular insufficiency or portal hypertension.

Hematological and biochemical examinations were normal (Table 1). There was no cholestasis or cytolysis. The levels of alpha-feto-protein and carcinoembryonic antigen were normal. The cytobacteriological examination of the urine was sterile.

**Table-I**  
*Hematological and biochemical parameters*

Parameters	Results
Hemoglobin	12.3 (g/dL)
Hematocrit	37,3 (%)
Leukocytes	4950 (/mm <sup>3</sup> )
Platelets	191 (/mm <sup>3</sup> )
Prothrombin activity	100 (%)
C-reactive protein	1,6 (mg/L)
AST	21 (UI/L)
ALT	12 (UI/L)
GGT	21 (U/L)
Alkaline phosphatase	67 (U/L)
Total bilirubin	6 (mg/L)
Albumin	46.8 (g/L)
$\alpha$ -Fetoprotein	1.6 (ng/mL)
CEA	3 (ng/mL)
LDH	349 (U/L)
Urea	0.35 (g/L)
Creatinine	6 (mg/L)

AST: aspartate aminotransferase

ALT: alanine aminotransferase

GGT : $\alpha$ -glutamyl transferase


CEA : carcinoembryonic antigen

LDH : lactate dehydrogenase

1. Gastroenterology Department, Mohammed VI University Hospital, Marrakech

2. Physiology Department, Faculty of Medicine and Pharmacy at Cadi Ayyad University, Marrakech

**Address of Correspondence:** Dr. Lemfadli Y, Gastroenterology Department, Mohammed VI University Hospital, Marrakech



**Figure 1:** Magnetic resonance imaging showing ectasias of the intrahepatic bile ducts and communication with the bile branches

Abdominal ultrasound showed cystic formations throughout the hepatic parenchyma which seem to communicate with the intrahepatic bile ducts. There is no calculus or liver nodule. Magnetic resonance imaging (MRI) (Figure 1) confirmed the diagnosis of CD. The liver was of normal size, with regular contours with the presence of multiple cystic formations in hyposignal T1 and hypersignal T2, with thin walls. These cystic lesions are connected to and communicate with the intrahepatic bile ducts.

Liver biopsy was not performed because the procedure would bring more risks than benefits. Currently, the patient continues her regular clinical follow-up in consultation. She has no major clinical manifestations that affect her daily life. The prognosis for her illness is favorable at the moment, since the patient has no complications including no cholangitis and no signs of portal hypertension or malignancy.

### Discussion

Caroli's disease is a rare autosomal recessive inherited congenital disease. Rare cases of autosomal dominant transmission have been reported.<sup>4</sup> Its incidence is estimated at 1 per million of the population [5]. It is associated with incomplete and defective remodeling of the embryonic ductal plate.<sup>6,7</sup> It is usually diagnosed in childhood or adolescence<sup>4</sup>, but can be diagnosed in adulthood.<sup>8-10</sup> No sex predominance has been reported.<sup>11</sup>

The clinical symptomatology of CD is not specific. Patients may be asymptomatic or have symptoms such as jaundice, right upper quadrant pain or fever.<sup>12</sup> Intrahepatic ductal ectasia predisposes to the stagnation of bile, which can lead to the formation of stones and predisposes to repeated cholangitis which can be complicated by serious infections such as liver abscess or septicemia.<sup>13,14</sup> Chronic inflammation due

to gallstones in the intrahepatic ducts can lead to secondary biliary cirrhosis with its complications such as portal hypertension.<sup>14</sup>

Liver function tests in CD may be normal or there may be reversible increases during episodes of cholangitis [15]. The level of alkaline phosphatase and direct bilirubin may be increased. The level of transaminases can be normal as it can be high and reflect a progressive hepatic fibrosis in case of Caroli syndrome.<sup>14</sup>

The final diagnosis is confirmed by histopathology, but current non-invasive imaging tools, including ultrasound (US), computed tomography (CT) and magnetic resonance imaging (MRI) are considered to be the first-line diagnostic modalities.<sup>16</sup> The ultrasound shows saccular or spindle-shaped cystic dilations predominant around the hepatic hilum without any underlying obstacle and which communicate with the rest of the biliary tree. It is a very good technique for *visualizing* intrahepatic lithiasis. Doppler ultrasound or multi-slice CT also enable visualization of the pathognomonic “dot sign”, which represents a portal or arterial branch at the periphery of or within a pseudoseptation running through a cyst.<sup>17</sup> Magnetic resonance imaging (MRI) is a non-invasive technique that currently presents the most specific tool for diagnosing CD. It demonstrates cystic structures of variable size which communicate freely with the biliary tree as well as the “dot sign” and intrahepatic stones.<sup>18</sup> Caroli syndrome differs from CD by the presence of smaller cystic structures (<2 cm) connected to the bile ducts with periportal fibrosis.<sup>19</sup>

The management of CD depends on the symptomatology of the patient and the extent of biliary abnormalities.<sup>10</sup> Cholangitis can be managed with antibiotics. In the case of cholestasis with intrahepatic lithiasis, the prescription of ursodeoxycholic acid is an option.<sup>17,20</sup> Surgical resection by lobectomy is recommended in case of localized disease in a lobe.<sup>21</sup> It reduces the risk of degeneration into cholangiocarcinoma which to be as high as 100-fold greater in patients with CD than in the general population.<sup>22</sup> However, in cases of diffuse disease and in case of suspected malignant transformation, liver transplantation remains the treatment of choice.<sup>23</sup> In case of biliary obstruction, an endoscopic sphincterotomy, radiological or surgical drainage may be applied.<sup>15</sup>

### Conclusion

Caroli disease poses a double diagnostic and therapeutic problem. The rarity of the disease, the absence of specific clinical signs and the frequent association with gallstones make diagnosis difficult.

Treatment of diffuse forms is problematic and requires liver transplantation given the risk of septic complications, secondary biliary cirrhosis and degeneration.

### References

1. Pimentel F. Enfermedad de Caroli. Rev Chil Cir 2004;56:426-433.
2. Gupta AK, Gupta A, Bhardwaj VK, Chansoria M. Caroli's disease. Indian J Pediatr 2006;73:233-235. <https://doi.org/10.1007/BF02825490> PMID:16567920
3. Menon S, Holt A. Large-duct cholangiopathies: aetiology, diagnosis and treatment. Frontline Gastroenterol. 2019;10(3):284-291. <https://doi.org/10.1136/flgastro-2018-101098> PMID:31288256 PMID:PMC6583582
4. R. Torra, C. Badenas, A. Darnell, C. Br 'u, A. Escorsell, and X. Estivill, “Autosomal dominant polycystic kidney disease with anticipation and Caroli's disease associated with a PKD1 mutation rapid communication,” *Kidney International*, vol.52, no. 1, pp. 33-38, 1997. <https://doi.org/10.1038/ki.1997.300> PMID:9211343
5. Bakoyiannis A, Delis S, Triantopoulou C, Dervenis C: Rare cystic liver lesions: a diagnostic and managing challenge. *World J Gastroenterol* 2013; 19(43):7603-7619. <https://doi.org/10.3748/wjg.v19.i43.7603> PMID:24282350 PMID:PMC3837259
6. Khalefa, A.A., Alrasheed, M. & Saeedan, M.B. Central dot sign. *Abdom Radiol* 41, 2289-2290 (2016). <https://doi.org/10.1007/s00261-016-0836-2> PMID:27392743
7. Sato Y, Harada K, Kizawa K, Sanzen T, Furubo S, Yasoshima M et al. (2005) Activation of the MEK5/ERK5 cascade is responsible for biliary dysgenesis in a rat model of Caroli's disease. *Am J Pathol* 166:49-60. [https://doi.org/10.1016/S0002-9440\(10\)62231-6](https://doi.org/10.1016/S0002-9440(10)62231-6)
8. Acioli, Maira Lima et al. “Diffuse Caroli's disease with atypical presentation: a case report.” *Annals of gastroenterology* vol. 27,1 (2014): 79-81.
9. Sarker, I., Bhuiyan, A. M., & Sarker, M. Z. (2016). A middle aged man with Caroli's disease: A case report. *Bangabandhu Sheikh Mujib Medical University Journal*, 9(1), 50-52. <https://doi.org/10.3329/bsmmuj.v9i1.28945>
10. Wang ZX, Li YG, Wang RL, Li YW, Li ZY, Wang LF, Yang HY, Zhu Y, Wang Y, Bai YF et al: Clinical classification of Caroli's disease: an analysis of 30 patients. *HPB (Oxford)* 2015; 17(3):278-283. <https://doi.org/10.1111/hpb.12330> PMID:25327281 PMID:PMC4333791
11. Madjov R, Chervenkov P, Madjova V, Balev B. (2005) Caroli's disease. Report of 5 cases and review of literature. *Hepatogastroenterology* 52:606-609.

12. Huang, Xiao-ming. "Factors contributing to diagnostic delay of Caroli syndrome: a single-center, retrospective study." (2020).
13. Cabral Correia P, Morgado B (September 20, 2017) Caroli's Disease as a Cause of Chronic Epigastric Abdominal Pain: Two Case Reports and a Brief Review of the Literature. *Cureus* 9(9): e1701. <https://doi.org/10.7759/cureus.1701> PMID:29159008 PMCID:PMC5690396
14. Yonem, Ozlem, and Yusuf Bayraktar. "Clinical characteristics of Caroli's disease." *World journal of gastroenterology* vol. 13,13 (2007): 1930-3. <https://doi.org/10.3748/wjg.v13.i13.1930> PMID:17461492 PMCID:PMC4146968
15. Yadav, Padmini et al. "Caroli's disease: a diagnostic challenge." *International Surgery Journal* 5 (2018): 3750. <https://doi.org/10.18203/2349-2902.isj20184657>
16. Cannella R, Giambelluca D, Diamarco M, Caruana G, Cutaia G, Midiri M, Salvaggio G: Congenital Cystic Lesions of the Bile Ducts: Imaging-Based Diagnosis. *Curr Probl Diagn Radiol* 2019. <https://doi.org/10.1067/j.cpradiol.2019.04.005> PMID:31027922
17. Caroli's disease: current knowledge of its biliary pathogenesis obtained from an orthologous rat model *Int J Hepatol*, 2012 (2012), pp. 1-10. <https://doi.org/10.1155/2012/107945> PMID:22007315 PMCID:PMC3168917
18. Guy F, Cagnet F, Dransart M, Cercueil JP, Conciatori L, Krause D. Caroli's disease: magnetic resonance imaging features. *Eur Radiol* 2002;12:2730-2736. <https://doi.org/10.1007/s00330-002-1471-6> PMID:12386765
19. Strazzabosco M, Fabris L, Spirli C. Pathophysiology of cholangiopathies. *J Clin Gastroenterol* 2005;39:S90-S102. <https://doi.org/10.1097/01.mcg.0000155549.29643.ad> PMID:15758666
20. Ananthakrishnan AN, Saeian K: Caroli's disease: identification and treatment strategy . *Curr Gastroenterol Rep.* 2007, 9:151-155. [10.1007/s11894-007-0010-7](https://doi.org/10.1007/s11894-007-0010-7) <https://doi.org/10.1007/s11894-007-0010-7> PMID:17418061
21. Chen CB, Hu WD, Zhao WW, Gu YY, Hou HW, Pan Z. Laparoscopic hepatectomy for the treatment of Caroli's disease: a case report. *Ann Surg Treat Res.* 2018 Mar;94(3):162-165. <https://doi.org/10.4174/ast.2018.94.3.162> PMID:29520352 PMCID:PMC5842090
22. Jang MH, Lee YJ, Kim H. Intrahepatic cholangiocarcinoma arising in Caroli's disease. *Clin Mol Hepatol* 2014;20:402-405. <https://doi.org/10.3350/cmh.2014.20.4.402> PMID:25548748 PMCID:PMC4278073
23. DeKerckhove L., De, Meyer M., Verbaandert, C., Mourad, M., Sokal, E., Goffette, P., Geubel, A., Karam, V., Adam, R., and Lerut, J. The place of liver transplantation in Caroli's disease and syndrome. *Transpl.Int.* 2006; 19: 381-388. <https://doi.org/10.1111/j.1432-2277.2006.00292.x> PMID:16623873.