



BJP

Bangladesh Journal of Pharmacology

Research Article

Antihyperlipidemic and renoprotective activities of methanolic extract of *Canscora decussata* extract in alloxan-induced diabetic rabbits

Antihyperlipidemic and renoprotective activities of methanolic extract of *Canscora decussata* extract in alloxan-induced diabetic rabbits

Nadeem Irshad^{1,2}, Muhammad Shoaib Akhtar², Yousaf Kamal¹, M. Imran Qayyum², Abdul Malik² and Hafiz Rashid Hussain²

¹Hamdard Institute of Pharmaceutical Sciences, Islamabad Campus, Islamabad 44000, Pakistan; ²Department of Pharmacology, Faculty of Pharmacy, University of Sargodha, Sargodha 40100, Pakistan.

Article Info

Received: 28 June 2013
Accepted: 13 July 2013
Available Online: 17 July 2013
DOI: 10.3329/bjp.v8i3.15506

Cite this article:

Irshad N, Akhtar MS, Kamal Y, Qayyum MI, Malik A, Hussain HR. Antihyperlipidemic and renoprotective activities of methanolic extract of *Canscora decussata* extract in alloxan-induced diabetic rabbits. Bangladesh J Pharmacol. 2013; 8: 323-27.

Abstract

The current study was designed to evaluate the hypolipidemic and renoprotective effects of methanolic extract of powdered *Canscora decussata* whole plant in the diabetic rabbits. Thirty rabbits were divided into five groups having 6 animals each including normal and diabetic controls groups, the remaining groups received methanolic extract in 400 and 600 mg/kg doses and another group got pioglitazone (3 mg/kg) for 30 days. Serum levels of triglycerides, total cholesterol, LDL-cholesterol, HDL-cholesterol, albumin, globulin and total proteins were estimated by using commercially available kits. The results showed that extract significantly ($p < 0.01$) decreased the raised parameters including triglyceride, total cholesterol and LDL-cholesterol, atherogenic index, Coronary risk index up to normal values compared to diabetic rabbits. However, it significantly increased HDL-cholesterol, albumin, globulin and total protein levels. Therefore, it is suggested that methanolic extract of *C. decussata* exerts hypolipidemic and renoprotective effects in the alloxan-induced diabetic rabbits.

Introduction

Hypercholesterolemia and hypertriglyceridemia have been reported to occur in the diabetic rabbits (Maciejewski et al., 2001; Wojtowicz et al., 2004). High density lipoproteins also decrease in type 2 diabetic patients which ultimately lead to atheromatous disease (Rang et al., 2003). The development of diabetic nephropathy has been characterized by progressive increase in the excretion of protein, particularly albumin leading to end stage renal failure (Alberti et al., 1997). Diabetic nephropathy has been considered a major cause of end-stage renal disease in many developed countries (CDC, 2005).

In developing countries like Pakistan, 80% of population depending on traditional medicine in primary health care (Grover and Yadav, 2004) because these

drugs are of low cost and free from adverse effects (Akhtar et al., 2011). Ethnopharmacological and ethno botanical surveys indicate that more than 1200 plants are used worldwide in herbal medicine to treat diabetes (Ziyyat et al., 1997; El-Hilaly et al., 2003; Tahraoui et al., 2007). Among them *Actinodaphne hookeri*, *Aegle marmelos*, *Bombax ceiba*, *Cajanus scarabaeoides*, *Eulophia herbacea*, *Gymnema lactiferum*, *Lagerstroemia speciosa*, *Mangifera indica*, *Meyna spinosa*, *Nigella sativa*, *Teucrium stocksianum* *Trichosanthes tricuspidata* have shown antidiabetic effect in their scientifically designed studies (Prajapati et al., 2008; Ravi et al., 2009; Pattanayak et al., 2009; Tatiya et al., 2013; Bandara et al., 2009; Saha et al., 2009; Bhowmik et al., 2009; Sen et al., 2013; Khanam et al., 2009; Alamgeer et al., 2013; Bhavsar and Talele, 2013; Kulandaivel et al., 2013).



The *Canscora decussata* plant (Family: Gentianaceae), locally known as Sankhahulee, has been used in the traditional medicine for the treatment of insanity, epilepsy and nervous debility. It has been found to contain triterpenes, alkaloids, and xanthenes (Kokate et al., 2002). Recently, Akhtar et al., 2012 have reported its hypoglycemic effects in normal as well as in diabetic rabbits. In folklore, it is reputed as a nervine tonic, alternative, and laxative (Shah et al., 2000). Diabetes affects both glucose and lipid metabolism (Sperling et al., 2000) while renal damage is its serious complication (Alberti, 1993).

Ipomoea staphylina leaves (Bag and Mumtaz, 2013), *Sechium eduleon* (Mumtaz et al., 2013), *Nigella sativaon* (Begum et al., 2006), *Cinnamomum zeylanicum* (Ullah et al., 2013), and grape seed and skin (Hamlaoui et al., 2012) extract shows renoprotective effect.

However, as far as ascertained, no scientific study was reported to assess the antihyperlipidemic and renoprotective activities of *C. decussata* in diabetes. Therefore, the current study was conducted to determine the antihyperlipidemic and renoprotective effects of methanolic extract of *C. decussata* in the diabetic rabbits.

Material and Methods

Chemicals and plant material

The plant *C. decussata* was collected from a village of Lahore, Pakistan in the month of August 2010 and got identified by the taxonomist of University of Sargodha, Sargodha. A voucher specimen (No. UOS/CD/333) was deposited in Department of Pharmacy, University of Sargodha. The plant was completely shade dried and powdered with herbal grinder. The powdered material was stored in well closed cellophane bags at 4°C in the refrigerator till further use. Chemicals used in the study include triglyceride, total cholesterol, LDL-cholesterol and HDL-cholesterol kits by Fluitest, Germany; alloxan monohydrate by Research Organics, USA; Methanol by Merck Chemical Co., Germany; and Gum acacia by Uni-Chem, Germany, pioglitazone was a generous gift from Maan Gee distributors, Sargodha.

Experimental animals

Healthy adult rabbits weighing 1,000-1,500 g were kept at animal house of the Department of Pharmacy, Hamdard University Islamabad Campus, Islamabad. The animals were housed in stainless cages under standard laboratory condition (light period: 8:00 am to 8:00 pm, 21 ± 2°C, relative humidity 55%, green fodder and water were available *ad libitum*).

Preparation of *C. decussata* extract

Methanolic extract of *C. decussata* whole plant was prepared by cold maceration. Then extract was dried with rotary evaporator.

Induction of diabetes

After overnight fasting, rabbits were made diabetic by

intravenous injection of fresh solution of 150 mg per kg of alloxan monohydrate in jugular vein. Rabbits received alloxan monohydrate was provided with a free access to water with 5% dextrose solution in order to protect them from hypoglycaemic shock (Akhtar et al., 2002). Three days (72 hours) after injecting the alloxan blood glucose level of surviving rabbits were measured with glucometer and rabbits having blood glucose level between 250-300 mg/dL were considered diabetic and used for further study (Olajide et al., 1999; Shani et al., 1974).

Administration of drug suspensions

The quantity of extract was calculated on body weight and triturated with about 10 mL of 2% aqueous gum solution and the final volume was always made up to 20 mL. Then suspension was administered (p.o) to each animal by using a stomach tube and disposable syringe (Akhtar and Iqbal, 1991; Sivajothi et al., 2008).

Experimental design

The rabbits were divided into five groups of six animals each. Group 1 and 2 considered as normal and diabetic control and were administered orally 20 mL of 2% aqueous gum acacia solution daily for 30 days. The groups 3 and 4 were administered extract (400 and 600 mg/kg OD), while group 5 received pioglitazone continuously for 30 days respectively. After 30 days blood samples were collected for the study.

Biochemical analysis

The total serum cholesterol, triglyceride and HDLs were estimated by enzymatic test kits using chemistry analyzer biolyser 100 (Merck Chemical Co., Germany). VLDL cholesterol and LDL-cholesterol were calculated using the Friedewald's et al., 1972 formula. Atherogenic index and coronary risk index was calculated by using Devaki et al., 2011 formula. Serum total protein and albumin content was determined by standard procedures in an auto analyzer using standard kits.

Statistical analysis

The results are expressed as mean ± SEM. The statistical analysis was carried out using paired t-test and one-way analysis (ANOVA). Statistical p value <0.05 was considered to be significant.

Results and Discussion

Effects of methanolic extract of *C. decussata* whole plant on serum triglyceride, cholesterol, LDL cholesterol, VLDL cholesterol, HDL-cholesterol, AI and CRI (mg/dL; mean ± SEM) levels in alloxan-induced diabetic rabbits after 30 days of extract administration are given in Table I. The serum levels of triglyceride (99.8 ± 0.31 mg/dL) and total cholesterol (129.9 ± 0.98 mg/dL) of group treated with 400 mg/kg extract were found to be significantly (p<0.001) lowered as compared to untreated alloxan-induced diabetic. Whereas, the serum level of HDL-cholesterol (34.6 ± 0.6 mg/dL) of group treated with 400 mg/kg extract was found to be

significantly ($p < 0.001$) increased as compared to untreated alloxan-induced diabetic. The AI and CRI also decreased significantly ($p < 0.001$) as compared to diabetic group. The serum levels of triglyceride (83.6 ± 0.7 mg/dL), total cholesterol (123.7 ± 0.78 mg/dL), LDL-cholesterol (67.3 ± 1.4 mg/dL) and VLDL-cholesterol (16.7 ± 0.4 mg/dL) of group treated with 600 mg/kg extract were found to be significantly ($p < 0.001$) lowered as compared to untreated alloxan-induced diabetic group. Whereas, the serum level of HDL-cholesterol (39.7 ± 1.1 mg/dL) of group treated with 600 mg/kg of methanolic extract were found to be significantly ($p < 0.001$) increased as compared to untreated alloxan-induced diabetic group. AI and CRI also significantly lowered as compared to diabetic control group. The serum level of HDL-cholesterol (37.4 ± 0.2 mg/dL) of group treated with pioglitazone were found to be significantly ($p < 0.001$) increased as compared to untreated alloxan-induced diabetic group (Table I).

Effects of methanolic extract of *C. decussata* whole plant on serum total protein, albumin, and globulin and A/G ratio in alloxan induced diabetic rabbits after 30 days of extract administration are given in Table II. The serum levels of total protein (7.5 ± 0.02 g/dL), albumin (4.5 ± 0.03 g/dL) of group treated with methanolic extract in dosing 400 mg/kg were found to be significantly ($p < 0.05$) increased as compared to untreated alloxan-induced diabetic group. The serum levels of total

protein (7.6 ± 0.02 g/dL), albumin (4.8 ± 0.03 g/dL), globulin (3.1 ± 0.3 g/dL) and A/G ratio (1.5 ± 0.1 g/dL) of group treated with extract in dosing 600 mg/kg were found to be significantly ($p < 0.05$) increased as compared to untreated alloxan-induced diabetic group. Similarly, the serum levels of total protein (6.5 ± 0.2 g/dL), albumin (4.2 ± 0.1 g/dL), globulin (2.3 ± 0.1 g/dL) and A/G ratio (1.8 ± 0.01 g/dL) of group treated with pioglitazone in dosing 3 mg/kg were found to be significantly ($p < 0.05$) increased as compared to untreated alloxan-induced diabetic group.

Discussion

Diabetic patients have more chances to develop atheromatous complications such as ischemic heart diseases (Way et al., 2001). The decrease level of high density lipoprotein in diabetic patient is also the leading cause of atheromatous diseases (Rang et al., 2003). Oral administration of methanolic extract of *C. decussata* cause an increase in HDL and reduction in total cholesterol, triglycerides, LDL and VLDL which protect diabetic patients from atheromatous disease. The continuous administration of extract produced beneficial effect against hyperlipidemia associated with hyperglycemia in a dose dependent manner. Standard drug pioglitazone also increased serum HDL while reduces the total cholesterol, triglycerides, LDL and VLDL in dose dependent manner as reported by

Table I

Effects of methanolic extract of *C. decussata* on serum cholesterol, triglyceride, LDL, VLDL, HDL, atherogenic index and coronary risk index (mg/dL) levels in alloxan-induced diabetic rabbits after 30 days of administration

Groups	Cholesterol mg/dL	Triglyceride mg/dL	HDL mg/dL	LDL mg/dL	VLDL mg/dL	Atherogenic index mg/dL	Coronary risk index mg/dL
Normal control	121.0 ± 0.9	80.6 ± 0.7	42.0 ± 1.2	58.7 ± 0.6	20.5 ± 0.7	1.3 ± 0.01	2.8 ± 0.02
Diabetic control	266.3 ± 1.5	218.8 ± 1.4	29.7 ± 0.7	191.3 ± 0.7	45.5 ± 1.0	6.4 ± 0.4	8.9 ± 0.3
Methanolic extract 400 mg	129.9 ± 1.0 ^b	99.8 ± 0.3 ^b	34.6 ± 0.6 ^b	75.3 ± 1.3 ^b	19.9 ± 0.6 ^b	2.1 ± 0.1 ^b	3.7 ± 0.2 ^b
Methanolic extract 600 mg	123.7 ± 0.8 ^b	83.6 ± 0.7 ^b	39.7 ± 1.1 ^b	67.3 ± 1.4 ^b	16.7 ± 0.4 ^b	1.6 ± 0.02 ^b	3.1 ± 0.3 ^b
Pioglitazone treated	147.5 ± 1.0 ^b	87.0 ± 0.03 ^b	37.4 ± 0.2 ^b	92.6 ± 1.2 ^b	17.4 ± 0.4 ^b	2.4 ± 0.01 ^b	3.9 ± 0.1 ^b

Values are expressed as mean ± SEM; n= 6; ^ap < 0.05, ^bp < 0.001, ^cp > 0.05 considered statistically significant as compared to diabetic control group

Table II

Effect of methanolic extract of *C. decussata* on the concentration of total protein, albumin, globulin and A/G ratio (g/dL) in serum of control and experimental groups after 30 days

Groups	Total protein	Albumin	Globulin	A/G Ratio
Normal control	8.0 ± 0.1	5.0 ± 0.02	3.1 ± 0.1	1.5 ± 0.03
Diabetic control	4.4 ± 0.03	2.3 ± 0.1	2.1 ± 0.1	1.0 ± 0.1
Methanolic extract 400 mg	7.5 ± 0.02 ^b	4.5 ± 0.03 ^a	2.7 ± 0.01 ^c	1.6 ± 0.04 ^a
Methanolic extract 600 mg	7.6 ± 0.02 ^b	4.8 ± 0.03 ^b	3.1 ± 0.3 ^a	1.5 ± 0.1 ^a
Pioglitazone treated	6.5 ± 0.2 ^a	4.2 ± 0.1 ^a	2.3 ± 0.1 ^a	1.8 ± 0.01 ^a

Values are expressed as mean ± SEM; n= 6; ^ap < 0.05, ^bp < 0.001, ^cp > 0.05 considered statistically significant as compared to diabetic control group

Kakadiya and Shah (2010). The antihyperlipidemic activity of the test extract would be due to its control of hyperglycemia which elevates the triglyceride, total cholesterol and LDL levels (Maciejewski et al., 2001). Several studies have reported that atherogenic index is an excellent predictor of HD risk and monitor for the effectiveness of lipid-lowering therapies. But, the LDL-C/HDL-C ratio has been considered more prognostic than LDL-C or HDL-C alone (Natarajan et al., 2003). In the present study the test extract significantly reduced the AI as well as TC/HDL-cholesterol ratio which show its protective effect against cardiovascular dis-eases. Similar results have reported for *B. tomentosa* which significantly reduced the atherogenic and cardiac risk index in diabetic rats (Devaki et al., 2011). The protein profile parameters are also disturbed during diabetes mellitus as reported by Sumana and Suryawanshi (2001) in rats. The extract of *C. decussata* has also shown improvement in the insulin secretion and reverses the altered protein profile in a dose dependent manner by exerting the protein sparing effect. Generally, significant increase in urinary excretion of protein, albumin and glucose levels indicates the impaired renal function in diabetes. However, the treatments with herbal remedies have been found to prevent such changes (Tedong et al., 2006). Peroxisome proliferator-activated receptor-gamma agonists have been reported to have direct beneficial effects on the diabetic renal diseases. They have shown to reduce proteinuria and improve glomerulosclerosis, both in animal and human diabetic nephropathic studies (Baylis et al., 2003), which is also shown in this study. In addition, Ohga et al., (2007) reported that pioglitazone ameliorates renal injury may be by the inhibition of NF- κ B activation, ICAM-1 expression and macrophage infiltration in streptozotocin-induced diabetic rats. The present study has thus duly supported the alleged medicinal use of the plant in the traditional medicine.

In conclusion methanolic extract of *C. decussata* has been observed to exert significant and consistent hypolipidemic and renoprotective effects in the diabetic rabbits which also confirm the presence of such active principle (s) which have the maximum solubility in methanol solvent.

References

- Akhtar MS, Iqbal J. Evaluation of hypoglycemic effect of *Achyranthes aspera*, Linn. In normal and diabetic rabbits. J Ethnopharmacol. 1991; 31: 49-57.
- Akhtar MS, Khan MA, Malik MT. Hypoglycemic activity of *Alpinia galangal* rhizome and its extracts in rabbits. Fitoterapia 2002; 73: 623-28.
- Akhtar MS, Nadeem M, Rashid, Bashir S. Hypoglycaemic activity of different fractions of *Berberis aristata* root-bark in normal and alloxan diabetic rabbits. Can J App Sci. 2011; 2: 16-28.
- Alamgeer, Rashid M, Bashir S, Mushtaq MN, Khan HU, Malik MNH, Qayyum A, Shafeeq ur Rahaman M. Comparative hypoglycemic activity of different extracts of *Teucrium stocksianum* in diabetic rabbits. Bangladesh J Pharmacol. 2013; 8: 186-93.
- Alberti KGMM, Zimmet P, Defronzo RA, Keen H. International textbook of diabetes mellitus. 2nd ed. Chichester, John Wiley and Sons, 1997, p 1379.
- Alberti KGMM, Problems related to definitions and epidemiology of type 2 diabetes mellitus: Studies throughout the world. Diabetologia, 1993; 36; 978-84.
- Bag AK, Mumtaz SMF. Hepatoprotective and nephroprotective activity of hydroalcoholic extract of *Ipomoea staphylina* leaves. Bangladesh J Pharmacol. 2013; 8: 263-68.
- Bandara T, Rokeya B, Khan S, Ali L, Ekanayake S, Jansz ER, Balasubramanium K. Effects of *Gymnema lactiferum* leaves on glycemic and lipidemic status in type 2 diabetes subjects. Bangladesh J Pharmacol. 2009; 4: 92-95.
- Batteridge J. Lipid disorders in diabetes mellitus. In: Textbook of diabetes. William G (eds). 2nd ed. London, Blackwell Science, 1997, pp 1-31.
- Bayle C, Atzpodien EA, Freshour G. Peroxisome proliferators activated receptor- γ agonist provides superior renal protection versus angiotensin-converting enzyme inhibition in a rat model of type 2 diabetes with obesity. J Pharmacol Exp Ther. 2003; 307: 854-60.
- Begum NA, Dewan ZF, Nahar N, Mamun MIR. Effect of n-hexane extract of *Nigella sativa* on gentamicin-induced nephrotoxicity in rats. Bangladesh J Pharmacol. 2006; 1: 16-20.
- Bhavsar C, Talele GS. Potential anti-diabetic activity of *Bombax ceiba*. Bangladesh J Pharmacol. 2013; 8: 102-06.
- Bhowmik A, Khan LA, Akhter M, Rokeya B. Studies on the antidiabetic effects of *Mangifera indica* stem-barks and leaves on non diabetic, type 1 and 2 diabetic model rats. Bangladesh J Pharmacol. 2009; 4: 110-14.
- Bnouham M, Mekhfi H, Legssyer A, Ziyat A. Medicinal plants used in the treatment of diabetes in Morocco. Int J Diabetes Metab. 2002; 10: 33-50.
- Buckingham RE, Al-Barazanji KA, Toseland CN. Peroxisome proliferators-activated receptor- γ agonist, rosiglitazone, protects against nephropathy and pancreatic islet abnormalities in Zucker fatty rats. Diabetes 1998; 47: 1326-34.
- Centers for Disease Control and Prevention (CDC). Incidence of end-stage renal disease among persons with diabetes-United States, 1990-2002. MMWR Morb Mortal Wkly Rep. 2005; 54: 1097-1100.
- Devaki K, Beulah U, Akila G, Narmadha R, Gopalakrishnan VK. Glucose lowering effect of aqueous extract of *Bauhinia tomentosa* L. on alloxan induced type 2 diabetes mellitus in Wistar albino rats. JBCP. 2011; 2: 167-74.
- El-Hilaly J, Hmamouchi M, Lyoussi B. Ethnobotanical studies and economic evaluation of medicinal plants in Taounate province (Northern Morocco). J Ethnopharmacol. 2003; 86: 149-58.
- Friedewald WT, Levy RI, Fradrickson DS. Estimation of concentration of low density lipoprotein cholesterol in plasma without the use of preparative ultracentrifuge. Clin Chem. 1972; 18: 499-502.
- Grover JK, Yadav SP. Pharmacological actions and potential uses of *Momordica charantia*: A review. J Ethnopharmacol, 2004; 93: 123-32.
- Hamlaoui S, Samti O, Mokni M, Limam N, Carrier A, Limam

- F, Amri M, Aouani E, Marzouki L. Protective effect of grape seed and skin extract on high dosage garlic-induced renal oxidative stress. *Bangladesh J Pharmacol.* 2012; 7: 54-62.
- Irshad N, Akhtar MS, Bashir S, Malik A, Kamal, Y. Hypoglycaemic effects of *Canscora decussata* (Schult) whole-plant in normal and alloxan-induced diabetic rabbits. *J Ethnopharmacol.* 2012; (under consideration)
- Kakadiya J, Shah NJ. Hypoglycemic and hypolipidemic activity of pioglitazone in normal and streptozotocin-nicotinamide induced diabetic rats. *Pharmacologyonline* 2010; 1: 1-7.
- Khanam M, Dewan ZF. Effects of the crude and the n-hexane extract of *Nigella sativa* Linn. (kalajira) upon diabetic rats. *Bangladesh J Pharmacol.* 2009; 4: 17-20.
- Kokate CK, Purohit AP, Gokhale SB. *Pharmacognosy* 19th ed. Pune, Nirali Prakashan, 2002; pp 524-25.
- Kulandaivel S, Bajpai P, Sivakumar T. Anti-hyperglycemic activity of *Trichosanthes tricuspidata* root extract. *Bangladesh J Pharmacol.* 2013; 8: 305-10.
- Laakso M. Epidemiology of diabetic dyslipidemia. *Diabetic Rev;* 1995; 3: 408-22.
- Maciejewski R, Rucinski P, Burski K, Figura T. Changes in glucose, cholesterol and serum lipid fraction levels in experimental diabetes. *Ann Univ Mariae Curie Sklodowska.* 2001; 56: 363-68.
- McCarthy KJ, Routh RE, Shaw W. Troglitazone halts diabetic glomerulosclerosis by blockade of mesangial expansion. *Kidney Int.* 2000; 58: 2341-50.
- Mumtaz SMF, Paul S, Bag AK. Effect of *Sechium edule* on chemical induced kidney damage in experimental animals. *Bangladesh J Pharmacol.* 2013; 8: 28-35.
- Natarajan S, Glick H, Criqui M, Horowitz D, Lipsitz SR, Kinoshian B. Cholesterol measures to identify and treat individuals at risk for coronary heart disease. *Am J Prev Med.* 2003; 25: 50-57.
- Ohga S, Shikata K, Yozai K. Thiazolidinedione ameliorates renal injury in experimental diabetic rats through anti-inflammatory effects mediated by inhibition of NF-κB activation. *Am J Physiol Renal Physiol.* 2007; 292: 1141-50.
- Olajide OA, Awe S, Makinde JM. Purgative effect of the methanol extract of *Morinda, Lucida*. *Fitoterapia* 1999; 70: 199-204.
- Pattanayak S, Nayak SS, Panda D, Shende V. Hypoglycemic of *Cajanus scarabaeoides* in glucose overloaded and streptozotocin-induced diabetic rats. *Bangladesh J Pharmacol.* 2009; 4: 131-35.
- Prajapati DD, Patel NM, Savadi RV, Akki KS, Mruthunjaya K. Alleviation of alloxan-induced diabetes and its complications in rats by *Actinodaphne hookeri* leaf extract. *Bangladesh J Pharmacol.* 2008; 3: 102-06.
- Ravi S, Sadashiva CT, Tamizmani T, Balasubramanian T, Rupeshkumar M, Balachandran I. *In vitro* glucose uptake by isolated rat hemi-diaphragm study of *Aegle marmelos* Correa root. *Bangladesh J Pharmacol.* 2009; 4: 65-68.
- Rahman AU, Zaman K. Medicinal plants with hypoglycemic activity. *J Ethnopharmacol.* 1989; 26: 1-55.
- Rang HP, Dale MM, Ritter JM, Moore PK (eds). *Atherosclerosis and lipoprotein metabolism.* In: The text-book of pharmacology. London, Churchill Livingstone/Elsevier, 2003, p 311.
- Roman-Ramos R, Florez-Saenz JL, Alarcon-Aguilar FJ. Antihyperglycemic effect of some edible plants. *J Ethnopharmacol.* 1995; 48: 25-32.
- Saha BK, Bhuiyan MNH, Mazumder Haque KKM. Hypoglycemic activity of *Lagerstroemia speciosa* L. extract on streptozotocin-induced diabetic rat: Underlying mechanism of action. *Bangladesh J Pharmacol.* 2009; 4: 79-83.
- Sen S, De B, Devanna N, Chakraborty R. Hypoglycaemic and hypolipidemic effect of *Meyna spinosa* leaves in high fat diet-alloxan induced type 2 diabetic rats. *Bangladesh J Pharmacol.* 2013; 8: 181-85.
- Shah Y, Tangalos EG, Petersen RC. Mild cognitive impairment. When is it a precursor to Alzheimer's disease. *Geriatrics* 2000; 55: 65-68.
- Shani J, Gold Schmied A, Joseph B, Abronson Z, Sneman FG. Hypoglycemic effects of *Trigonella faenum graecum* and *Lupinus termis* (Leguminosae) seed and their major alkaloids in alloxan-diabetic and normal rats. *Arch Int Pharmacodyn Ther.* 1974; 210: 27-37.
- Sivajothi V, Dey A, Jayakar B, Rajkapoor B. Antihyperglycemic and anti-oxidant effect of *Phyllanthus rheedii* on streptozotocin-induced diabetic rats. *Iranian J Pharma Res.* 2008; 7: 53-59.
- Sperling MA, Saunders PA. Diabetes mellitus. In: Nelson textbook of pediatrics. Behrman RE, Kliegman RM, Jenson HB (eds). 2000, pp 1767-91.
- Sumana G, Suryawanshi SA. Effect of *Vinca rosea* extracts in treatment of alloxan diabetes in male albino rats. *Ind J Exp Bio.* 2001; 39: 748-59.
- Tahraoui A, El-Hilaly J, Israili ZH, Lyoussi B. Ethnopharmacological survey of plants used in the traditional treatment of hypertension and diabetes in South-Eastern Morocco (Errachidia province). *J Ethnopharmacol.* 2007; 110: 105-17.
- Tatiya AU, Puranik PM, Surana SJ, Patil YS, Mutha RE. Evaluation of hypolipidemic, antidiabetic and anti-oxidant activity of *Eulophia herbacea* tubers. *Bangladesh J Pharmacol.* 2013; 8: 269-75.
- Tedong L, Dimo T, Dzeufi PD, Asongalem E, Sokeng DS, Callard P, Flejou F, Kamtchouing P. Antihyperglycemic and renal protective activities of *Anacardium occidentale* leaves in streptozotocin induced diabetic rats. *Afr J Trad CAM.* 2006; 3: 23-35.
- Ullah N, Khan MA, Khan T, Ahmad W. Bioactive traditional plant *Cinnamomum zeylanicum* successfully combat against nephrotoxic effects of aminoglycosides. *Bangladesh J Pharmacol.* 2013; 8: 15-21.
- Way KJ, Kata IN, King GL. Protein kinase C and the development of diabetic vascular complication. *Diabet Med.* 2001; 18: 945.
- Wojtowicz Z, Wrona W, Kis G, Blaszczyk M, Solecka A. Serum total cholesterol, triglyceride and high density lipoprotein levels in rabbits during the course of experimental diabetes. *Ann Univ Mariae Curie Sklodowska,* 2004; 59: 258-60.
- Ziyat A, Legssyer A, Mekhfi H, Dassouli A, Serhrouchni M, Benjelloun W. Phytotherapy of hypertension and diabetes in oriental Morocco. *J Ethnopharmacol.* 1997; 58: 45-54.

Author Info

Nadeem Irshad (Principal contact)

e-mail: ndmirshad@gmail.com