

BJP

Bangladesh Journal of Pharmacology

Research Article

Anti-oxidant and hepatoprotective effects of *Lithocarpus polystachyus* against carbon tetrachloride-induced injuries in rat

Anti-oxidant and hepatoprotective effects of *Lithocarpus polystachyus* against carbon tetrachloride-induced injuries in rat

Shenghua Li^{1,2,3}, Junying Zeng^{1,2,3}, Juan Tan^{1,2,3}, Jian Zhang^{1,2,3}, Qifan Wu^{1,2,3},
Lingpeng Wang^{1,2,3} and Xianjin Wu^{1,2,3}

¹The Department of Life Science, Huaihua University, Huaihua 418 008, China; ²Key Laboratory of Hunan Province for Study and Utilization of Ethnic Medicinal Plant Resources, Huaihua 418 008, China; ³Key Laboratory of Hunan Higher Education for Hunan-Western Medicinal Plant and Ethnobotany, Huaihua 418 008, China.

Article Info

Received: 25 September 2013
Accepted: 20 November 2013
Available Online: 3 January 2014
DOI: 10.3329/bjp.v8i4.16458

Cite this article:

Li S, Zeng J, Tan J, Zhang J, Wu Q, Wang L, Wu X. Anti-oxidant and hepatoprotective effects of *Lithocarpus polystachyus* against carbon tetrachloride-induced injuries in rat. Bangladesh J Pharmacol. 2013; 8: 420-27.

Abstract

The present study aims to investigate the hepatoprotective and anti-oxidant effects of the total flavonoid of *Lithocarpus polystachyus* Rehd.(LP-F) *in vitro* and *in vivo*. The *in vitro* anti-oxidant property of total flavonoids was investigated by employing various established systems. Rats with carbon tetrachloride-induced liver injury were used to assess the hepatoprotective and anti-oxidant effect of total flavonoids *in vivo*. The level of activity of glutamate pyruvate transaminase, aspartate aminotransferase, alkaline phosphatase and gamma-glutamyl transpeptidase, total bilirubin, total cholesterol, triglycerides total protein and albumin contents in the serum and malondialdehyde, superoxide dismutase, catalase, and glutathione in the liver and kidney of the rats were assayed using standard procedures. The results showed the total flavonoids of *L. polystachyus* has strong hepatoprotective and anti-oxidant activity *in vitro* and *in vivo*. These data were supplemented with histopathological studies of rat liver sections. This suggests that the hepatoprotective activity of formulation is possibly attributed to its free radical scavenging properties.

Introduction

The liver is a vital organ and has many important functions, including metabolism and detoxification of hepatotoxicants (Altasey et al., 2011). In most cases, liver damage is a widespread pathology which involve oxidative stress and a progressive evolution from steatosis to chronic hepatitis, fibrosis, cirrhosis, and hepatocellular carcinoma (Srivastava and Shivanandappa, 2010). Many people die every year due to liver anomalies. Among them, most common are cirrhosis, cholestasis, hepatitis, portal hypertension, hepatic encephalopathy, hepatic failure and certain tumors like hepatoma.

As one of the aspects of the body's natural ecosystem, it

is being realized that the majority of the diseases/disorders are mainly due to the hyper physiological burden of free radicals, causes imbalance in homeostatic phenomena between oxidants and anti-oxidants in the body. CCl₄ is a classical hepatotoxin leading to hepatic diseases, and is generally used in animal models for inducing acute liver injury (Weber, Boll and Stampfl, 2003). Reductive dehalogenation of CCl₄, by the P₄₅₀ enzyme system in the endoplasmic reticulum, into highly reactive trichloromethyl-free radicals ('CCl₃ or CCl₃OO') initiates the process of lipid peroxidation. This is considered to be the most important mechanism in the pathogenesis of liver damage induced by CCl₄ (Altas et al., 2011). CCl₄-induced damage is also able to change the anti-oxidant status of the tissues, which is



manifested by abnormal histopathological alteration with steatosis, centrilobular necrosis and cirrhosis in the liver (Xu et al., 2010).

Hepatoprotective and anti-oxidant effects have been investigated recently in different plant extracts due to their potent antioxidant activities, such as *Carum copticum* (Gilani et al., 2005), *Trianthema decandra* (Balamurugan and Muthusamy, 2008), *Anogeissus leiocarpus* (Atawodi et al., 2011), *Impatiens bicolor* (Nisar et al., 2012); *Meconopsis integrifolia* (Maxim.) Franch (Zhou et al., 2013), Huangshan Maofeng (Lu et al., 2013) and *Litchi chinensis* (Basu et al., 2012). However, many other plants, including *Lithocarpus polystachyus* have not been studied for such effects.

L. polystachyus of family Fagaceae, is a shrub distributed widely in the mountainous area in southern China (Institute of Botany, 1972). Its tender leaves have been a traditional Chinese herb and been taken as a sweet tonic drink for several hundreds of years (China Pharmacopoeia Committee, 1999). In the course of seeking the sweet components in the leaves, the sweet compounds were identified phlorizin, phloretin, quercetin, dihydrochalcone-2'- β -D-glucopyranoside, luteolin, and quercitrin (Li et al., 2010), and their content in the leaves was up to around 10.6% (Li et al., 2008), *L. polystachyus* is claimed to possess multiple bioactivities, including hepatoprotective, anti-hypertensive, antioxidant, vasodilative, cerebrovascular, cardiovascular, hypoglycemic, and antiobesity activities (Dong et al., 2012; Hou et al., 2012; Liao et al., 1994; Yang et al., 2007). According to the Dictionary of Chinese Traditional Drugs, hepatoprotective activity is the major pharmacological effect of *L. polystachyus* but no scientific data has been reported for the hepatoprotective effect of this medicinal plant so far.

The present study was carried out to investigate the hepatoprotective effect and a possible underlying antioxidant activity of indifferent extracts of the leaves of *L. polystachyus* by assaying various marker enzymes, antioxidant enzymes in carbon tetrachloride-induced rats.

Materials and Methods

Plant materials

L. polystachyus was collected from the Huaihua county of the western region of Hunan province, China in March, 2010 and authenticated by Prof. Xianjin Wu, Department of Bioscience, Huaihua University, where a voucher specimen has been deposited. The plant material was air dried indoor at room temperature before extraction.

Experimental animal

Male Wistar rats (220 \pm 4.5 g) purchased from the animal house of Xiangya School of Medicine of Central South University. They were kept in departmental

animal house in an environment-tally control room (25 \pm 2°C), relative humidity 50 \pm 5% RH and with 12 hours light and 12 hours darkness cycle. All the animals were free to standard laboratory feed and tap water before experiment. All the studies were performed in accordance with the guidelines for the care and use of laboratory animals, as adopted and promulgated by the Institutional Animal Care Committee.

Determination and preparation of total flavonoids of *L. polystachyus*

The extract from *L. polystachyus* was obtained by ultrasound-assisted extraction with 70% ethanol for 45 min (Li et al., 2008), and then filtrated through filter paper. The filtrate was evaporated till dryness under reduced pressure weighed, the yield of the extract 15.2%. The extract was solved with hot water, added into column loading-treated AB-8 macroreticular resin for adsorption for 12 hours, and then washed with 20% ethanol respectively for getting rid of impurity. LP-F in the column was eluted with 60% ethanol, eluting solution was evaporated till dryness under reduced pressure and weighed, the yield of total flavonoids of *L. polystachyus* 8.7%.

Preparation of standard curve: A calibration rutin solution containing 200.8 μ g/mL rutin was prepared with 70% ethanol. In all 0, 1, 2, 3, 4, 5 and 6 mL of calibration rutin solution was respectively added to seven 25 mL volumetric flasks; to each was added 6 mL of 70% ethanol and 1 mL 5% NaNO₂. Each flask was shaken fully, mixed and then stand still for 6 min; each flask was added 1 mL 10% AlCl₃ solution, shake them fully again, mixed and stand still 6 min subsequently, 10 mL 10% NaOH solution was added into each volumetric flask and dilution to the mark with 70% ethanol, shaken fully, mixed and then let it rest for 15 min, Its absorbency (A) was measured at 500 nm, 70% ethanol as a blank control.

One mL of above the extraction was transferred to a 25 mL volumetric flask, to measure its absorbance (A). Using the methodology described above, we finally calculated its total flavones content according to a regression equation

In vitro anti-oxidant activity

Scavenging activity on DPPH radical

The radical scavenging activity of total flavonoids of *L. polystachyus* was determined according to our previous procedure (He et al., 2012). In brief, 3.0 mL of 0.1 mM DPPH in methanol solution was added to 1.0 mL of total flavonoids of *L. polystachyus* preparative solution at different concentrations. The mixture was stood for 20 min in the dark before the absorbance was measured with the visible spectrometer at 517 nm. The DPPH scavenging effect was calculated as follows: DPPH scavenging activity (%) = [1 - (Absorbance of sample - Absorbance of background)/Absorbance of blank] \times

100.

Scavenging activity on ABTS radical

The ABTS radical cation scavenging activity of DDP was identified using a method (Re et al., 1999) with some modifications. ABTS was produced from the reaction of 7 mM of ABTS solution with 2.45 mM of potassium persulfate. The mixture was then kept in the dark at room temperature for 16 hours. At the time of use, the ABTS solution was diluted with ethanol to yield an absorbance of 0.70 ± 0.02 at 734 nm. Samples (0.2 mL) of various concentrations (0.2 to 1.0 mg/mL) were mixed with ABTS solutions (2 mL). After reaction at room temperature for 6 min, the absorbance at 734 nm was recorded. The ABTS scavenging effect was calculated based on the following formula: $\text{ABTS scavenging activity (\%)} = [1 - (\text{Absorbance of sample} - \text{Absorbance of background}) / \text{Absorbance of blank}] \times 100$.

Scavenging activity on hydroxyl radical

The HO⁻-scavenging activity of LP-F was measured by an improved Fenton-type reaction (Wang et al., 2012). In brief, 1.0 mL of LP-F, at different concentrations, was spiked with 1.0 mL of 6 mM FeSO₄, 1.0 mL of 6 mM salicylic acid-ethanol and 1.0 mL of 6 mM H₂O₂ and incubated at 37°C for 1 hour. BHT was used as positive control. The absorbance at 510 nm was measured with a visible spectrometer. The percentage of HO⁻-scavenging activity of total flavonoids of *L. polystachyus* was calculated according to the following formula: $\text{Scavenging activity against HO}^- (\%) = [1 - (\text{Absorbance of sample} - \text{Absorbance of blank}) / \text{Absorbance of control}] \times 100$.

Scavenging activity on superoxide radical

The scavenging activity on O₂^{•-} of total flavonoids of *L. polystachyus* was determined as described in our previous study (Wang et al., 2012). 1.0 mL of Mm nitroblue tetrazolium, 1.0 mL of mM nicotinamide adenine dinucleotide hydrogen, 1.0 mL of different concentration sample solution and 0.4 mL of 3 mM phenazine methosulfate were successively added into test tube, the reaction mixture was incubated at ambient temperature for 5 min and the absorbance of the mixture solution was determined at 560 nm with a visible spectrometer. O₂^{•-} scavenging capability of total flavonoids of *L. polystachyus* was calculated according to the following formula: $\text{Suppression rate (\%)} = \text{O}_2^{\bullet-} (\%) = [1 - (\text{Absorbance of sample} - \text{Absorbance of blank}) / \text{Absorbance of control}] \times 100$.

In vivo studies

Carbon tetrachloride-induced hepatotoxicity

Rats (n = 60) were divided into six groups consisting of 10 animals in each group. Group I served as normal group and received distilled water with 0.3% sodium carboxy-methylcellulose (CMC-Na) (1 mL/kg body

weight, p.o.) for 5 days, and olive oil (1 mL/kg bodyweight, s.c.) on days 2 and 3 (Jain et al., 2008). Group II served as control group and received 0.3% CMC-Na (1 mL/kg body weight, p.o.) for 5 days, and CCl₄-olive oil (1:1, 2 mL/kg body weight, s.c.) on days 2 and 3. Group III was treated with the standard drug silymarin (100 mg/kg body weight, p.o.) daily for 5 days, and also received CCl₄-olive oil (1:1, 2 mL/kg body weight, s.c.) on days 2 and 3, 30 min after administration of silymarin. Groups IV-VI (test group animals) were administered orally a dose of 100, 200, and 400 mg/kg body weight of total flavonoids of *L. polystachyus* (p.o.), respectively, for 5 days; in addition, they received CCl₄-olive oil (1:1, 2 mL/kg, s.c.) on days 2 and 3, 30 min after administration of total flavonoids of *L. polystachyus*. On day 6, animals were anesthetized, blood was collected and allowed to clot, and serum was separated for assessment of biochemical parameters. The rats were then sacrificed by bleeding, the liver and kidney were carefully dissected and cleaned of extraneous tissue, and part of the liver tissue was immediately transferred into 10% formalin for histopathological investigation.

Serum analysis for liver marker enzymes

Serum samples were assayed for alkaline phosphatase, glutamate pyruvate transaminase, aspartate aminotransferase, gamma-glutamyl transpeptidase by assay kits which were obtained from the Institute of Biological Engineering of Nanjing Jiancheng (Nanjing, China).

Serum analysis for biochemical studies

Total bilirubin, total cholesterol, triglycerides total protein and albumin were estimated by assay kits which were obtained from the Institute of Biological Engineering of Nanjing Jiancheng (Nanjing, China).

Determinations of antioxidant activity in vivo

The liver and kidney of superoxide dismutase, catalase, malondialdehyde and glutathione were determined by assay kits which were obtained from the Institute of Biological Engineering of Nanjing Jiancheng (Nanjing, China).

Histopathological studies

Liver tissues were fixed in 10% formalin for at least 24 hours, embedded in paraffin, and cut into 5 μm thick sections in a rotary microtome. The sections were stained with hematoxylin-eosin dye and observed under a microscope (IX51, Olympus, Japan) to detect histopathological changes in the liver.

Statistical analysis

Data obtained from this work were analyzed statistically using Students' t-test and ANOVA (One- or Two-way) followed by a post test (Tukey-Kramer multiple comparison test). Differences between means will be considered significant at 5% level of significance (p < 0.05).




Figure 1: *In vitro* free radical scavenging activity and antioxidant effects of LP-F. (A) DPPH-scavenging activity of various concentrations of total flavonoid of *L. polystachyus* and reference BHT. (B) ABTS-scavenging effect of LP-F and BHT. (C) HO[•]-scavenging activity of LP-F and BHT. (D) O₂^{•-}-scavenging activity of total flavonoid of *L. polystachyus* and BHT, Data are presented as means \pm SD (n = 3)

Results

Total flavonoids of *L. polystachyus* was extracted from *L. polystachyus* with 70% ethanol and purified with AB-8 macroporous resin, dried under reduced pressure and weighed, the yield of 8.7%. Regression equation: $A = 10.014C - 0.007$, $r = 0.9998$, C (concentration), with a better linear correlation between 7.9 and 54.4 µg/mL. Quantitative analysis indicated that the content of total flavonoids of *L. polystachyus* was up to 76.8%.

In this study, the *in vitro* antioxidant capacity of total flavonoids of *L. polystachyus* was estimated with DPPH, ABTS, hydroxyl radical, and reducing power systems. As shown in Figure 1A, total flavonoids of *L. polystachyus* concentration-dependently displayed DPPH-scavenging effects of 27.9, 39.9, 50.1, 62.4, 71.5, 79.2 and 84.7% at the tested concentrations of 25, 50, 75, 100, 125, 150 and 200 µg/mL, respectively. Similarly, total flavonoids of *L. polystachyus* at the concentrations of 25–200 µg/mL also exhibited obvious scavenging activity (31.9, 36.9, 53.6, 67.2, 76.2, 81.3 and 86.6%) against ABTS, respectively (Figure 1B). Total flavonoids of *L. polystachyus* also showed the significant scavenging

effects (49.5, 59.3, 70.3, 77.6, 82.5 and 86.6%) against HO[•] in a concentration-dependent manner (125–500 µg/mL, Figure 1C). As shown in Figure 1D, total flavonoids of *L. polystachyus* also showed the significant scavenging effects (24.6, 45.8, 55.5, 69.5, 79.6 and 85.6%) against O₂^{•-} in the broad range of 25–150 µg/mL. The data presented here indicates that total flavonoids of *L. polystachyus* has potential to be explored as a strong antioxidant, although it is inferior to the reference BHT in antioxidant capacity (Figure 1).

As depicted in Table I, the enzymatic activities of serum glutamate pyruvate transaminase, aspartate aminotransferase, alkaline phosphatase and gamma-glutamyl transpeptidase in CCl₄-intoxicated rats sharply increased to 135.2 ± 11.3 ($p < 0.001$), 223.5 ± 15.6 ($p < 0.001$), 197.5 ± 16.3 ($p < 0.001$) and 132.6 ± 5.8 IU/L ($p < 0.001$) from 32.6 ± 2.9 , 65.3 ± 1.7 , 48.3 ± 1.7 and 31.3 ± 2.2 IU/L of the normal group, respectively. Administration of LP-F (100, 200 and 400 mg/kg) alleviated the toxicity of CCl₄ and the changed serum level of glutamate pyruvate transaminase, aspartate aminotransferase, alkaline phosphatase and gamma-glutamyl

Table I

Protective effect of total flavonoid of *L. polystachyus* on liver marker enzymes

Group	Glutamate pyruvate transaminase (IU/L)	Aspartate aminotransferase (IU/L)	Alkaline phosphatase (IU/L)	Gamma-glutamyl transpeptidase (IU/L)
Normal	32.6 ± 2.9	65.3 ± 1.7	48.3 ± 1.7	31.3 ± 2.2
Control	135.2 ± 11.3 ^a	223.5 ± 15.6 ^a	197.5 ± 16.3 ^a	132.7 ± 5.8 ^a
Silymarin 100 (mg/kg)	48.7 ± 5.6 ^b	72.8 ± 4.9 ^b	61.4 ± 5.3 ^b	41.1 ± 4.2 ^b
Extract 100 (mg/kg)	105.7 ± 8.5 ^c	136.5 ± 19.6 ^c	135.4 ± 8.5 ^c	102.9 ± 7.6 ^c
Extract 200 (mg/kg)	68.5 ± 2.7 ^b	98.7 ± 8.5 ^b	109.4 ± 7.6 ^c	76.6 ± 4.4 ^b
Extract 400 (mg/kg)	54.3 ± 4.6 ^b	76.5 ± 6.3 ^b	76.2 ± 5.0 ^b	52.4 ± 6.5 ^b

Values are expressed as mean ± SEM, n = 10; ^ap<0.001 considered statistically significant as compared to normal control group; ^bp<0.001, ^cp<0.05 considered statistically significant as compared to carbon tetrachloride treated group

Table II

Protective effect of LP-F on serum biochemical profile

Group	Total bilirubin (mg/dL)	Total cholesterol (mg/dL)	Triglycerides (mg/dL)	Total protein (g/dL)	Albumin (g/dL)
Normal	0.6 ± 0.9	68.6 ± 5.6	32.7 ± 1.0	8.2 ± 0.7	1.0 ± 0.1
Control	1.9 ± 0.2 ^a	113.6 ± 6.6 ^a	79.4 ± 2.2 ^a	4.9 ± 0.1 ^a	2.7 ± 0.0 ^a
Silymarin 100 (mg/kg)	0.8 ± 0.1 ^b	73.7 ± 3.3 ^b	38.6 ± 4.2 ^b	7.6 ± 0.4 ^b	1.1 ± 0.1 ^b
Extract 100 (mg/kg)	1.4 ± 0.1 ^c	98.7 ± 4.7 ^c	71.6 ± 1.2 ^c	5.6 ± 0.3 ^c	2.1 ± 0.1
Extract 200 (mg/kg)	1.2 ± 0.1 ^c	87.7 ± 2.9 ^c	52.3 ± 1.0 ^c	7.0 ± 0.3 ^c	1.8 ± 0.2 ^c
Extract 400 (mg/kg)	0.8 ± 0.1 ^b	76.4 ± 7.7 ^b	40.6 ± 1.1 ^b	7.4 ± 0.9 ^b	1.2 ± 0.1 ^b

Values are expressed as mean ± SEM, n = 10; ^ap<0.001 considered statistically significant as compared to normal control group; ^bp<0.001, ^cp<0.05 considered statistically significant as compared to carbon tetrachloride treated group

transpeptidase, reverted towards the normal group with significance difference from the control group. Similarly, administration of silymarin (100 mg/kg) alleviated the toxicity of CCl₄ and serum level of glutamate pyruvate transaminase, aspartate aminotransferase, alkaline phosphatase and gamma-glutamyl transpeptidase was observed significantly less than that of control group.

The levels of serum total bilirubin, total cholesterol, triglycerides, total protein and albumin were estimated and the results were given in Table II. Administration of CCl₄ in rats caused a significant increase (p<0.05) in serum total bilirubin, total cholesterol, triglycerides and albumin levels, ranging from 0.6 ± 0.9, 68.6 ± 5.6, 32.6 ± 1.0 mg/dL, 8.2 ± 0.7 and 1.0 ± 0.1 g/dL for the normal control rats to 1.9 ± 0.2, 113.6 ± 6.6, 79.4 ± 2.2 mg/dL, 4.9 ± 0.1 and 2.7 ± 0.0 g/dL, while caused a significant decrease (p<0.05) in serum total protein ranging from 8.2 ± 0.7 g/dL for the normal rats to 4.9 ± 0.1 g/dL respectively. However, the protective administration of indifferent dose of LP-F decreased the CCl₄-elevated levels of serum total bilirubin, total cholesterol, triglycerides and albumin (p<0.05), and silymarin as positive control also exhibited a similar effect. While increased the CCl₄-elevated levels of serum total protein.

Determinations of the *in vivo* anti-oxidant activities of

total flavonoids of *L. polystachyus* in liver and kidney are presented in Figure 2. Compared with the normal group, the CCl₄-intoxicated rats exhibited a significant decrease in total superoxide dismutase, catalase and glutathione (liver and kidney) (p<0.001), significant increase in the level of malondialdehyde (in liver and kidney) (p<0.001). These changes were significantly reversed after treatment with LP-F and silymarin.

Histopathological observations of liver sections from the control group showed normal cellular architecture with distinct hepatic cells, sinusoidal spaces and a central vein (Figure 3A). In contrast, the CCl₄ group revealed the most severe damage of any of the groups; the liver sections showed massive fatty changes, necrosis, ballooning degeneration, broad infiltration of lymphocytes, and the loss of cellular boundaries (Figure 3B). The liver sections of the rats treated with total flavonoids of *L. polystachyus* (Figure 3E-F) showed a more or less normal lobular pattern with a mild degree of fatty change, necrosis, and lymphocyte infiltration almost comparable to the control (Figure 3A) and the silymarin-treated group (Figure 3C).

Discussion

The assays for scavenging DPPH, ABTS, HO⁻ and O₂⁻ showed that total flavonoids of *L. polystachyus* had a




Figure 2: Total flavonoid of *L. polystachyus* on liver and kidney superoxide dismutase, catalase, glutathione and malondialdehyde in CCl₄-intoxicated mice

Values are expressed as mean \pm SEM; n = 10; *p<0.001 considered statistically significant as compared to normal control group; ^bp<0.001, ^cp<0.05 considered statistically significant as compared to carbon tetrachloride treated group; Group I: Normal, Group II: control, Group III: silymarin 100 (mg/kg), Group IV: total flavonoid of *L. polystachyus* 100 (mg/kg), Group V: total flavonoid of *L. polystachyus* 200 (mg/kg), Group VI: total flavonoid of *L. polystachyus* 400 (mg/kg)

concentration-dependent anti-oxidant effect *in vitro*, and this effect was further verified by testing the anti-oxidant potential of total flavonoids of *L. polystachyus* in an animal model with experimental CCl₄-induced oxidative stress.

In this study, elevated levels of liver marker enzymes such as glutamate pyruvate transaminase, aspartate aminotransferase, alkaline phosphatase and gamma-glutamyl transpeptidase in serum are obtained with CCl₄ treatment indicating the pathophysiology of liver. Generally, CCl₄ is metabolized by the liver in to highly reactive metabolites which either directly or indirectly cause lipid peroxidation of the hepatocytes. Different cytosolic liver marker enzymes would then leaked out from these swollen and necrotic hepatocytes in to blood circulation and evidently elevated levels are obtained that is associated with the massive centrilobular necrosis, ballooning, degeneration and cellular infiltration of the liver (Singh et al., 2008). Treatment of rats by both CCl₄ and the LP-F/silymarin reversed the increase in liver marker enzymes towards the control group and hepatic lesions are also minimized (Table I).

It is well known that the liver play a pivotal role in the regulation of various chemicals. Administration of CCl₄ causes hepathopathy as indicated by elevation in serum level of total bilirubin, total cholesterol, triglycerides and albumin whereas decreases total protein in rat. These pathological changes signify the potential damage in hepatic induced with CCl₄ treatment. Treatment of rat with CCl₄ and with either LP-F or silymarin ameliorated the toxic effects of CCl₄ and restored the level of above biochemicals towards the normal group in accordance with other findings (Lin et al., 2008).

Carbon tetrachloride is accumulated in hepatic parenchyma cells and metabolized to the CCl₃[•] (Recknagel, 1983). CCl₃[•] radical reacts very rapidly with oxygen to yield a highly reactive CCl₃OO[•]. These radicals react with proteins and lipids. They remove hydrogen atoms from unsaturated lipids thus initiating lipid peroxidation, which causes loss of integrity of cell membranes and damage to hepatic tissue (Zhou et al., 2010). The enzymes superoxide dismutase, catalase and glutathione are key anti-oxidant enzymes that protect




Figure 3: Sections of the livers of CCl₄-treated rats showing the ventral vein (CV) and hepatic cells (hematoxylin-eosin staining, 200x) (A) Normal group, (B) Control group, (C) CCl₄ + silymarin (100 mg/kg), (D) CCl₄ + total flavonoid of *L. polystachyus* (100 mg/kg), (E) CCl₄ + total flavonoid of *L. polystachyus* (200 mg/kg), and (F) CCl₄ + total flavonoid of *L. polystachyus* (400 mg/kg).

against oxidative stress and tissue damage (Halliwell and Gutteridge, 1990).

These enzymes are critical for defence mechanisms against the harmful effects of reactive oxygen species (ROS) and free radicals in biological systems. The superoxide dismutase converts superoxide radicals into H₂O₂ and O₂, thus participating with other anti-oxidant enzymes, in the enzymatic defense against oxygen toxicity. Catalase is a key component of the anti-oxidant defense system. Inhibition of this protective mechanism results in enhanced sensitivity to free radical induced cellular damage. The decrease of catalase may result in a lot of deleterious effects due to the accumulation of superoxide radicals and hydrogen peroxide (Srinivasan et al., 2007). glutathione content was another important parameter that revealed oxidative damage in both liver and kidney. glutathione constitutes the first line of defense against free radicals. Reduction in liver glutathione activity in carbon tetrachloride-treated rats as observed in this study indicates the damage to the hepatic cells. Malondialdehyde is a cytotoxic product that is a hallmark of lipid peroxidation. The fact that LP-F treatment reduced elevated malondialdehyde and increased levels of superoxide dismutase, catalase and glutathione, indicated that it prevent the peroxidation of lipids by carbon tetrachloride.

Conclusion

L. polystachyus possessed strong hepato-protective and anti-oxidant activity in a rat model of CCl₄-induced.

The hepatoprotective activity may be due to its free radical-scavenging and anti-oxidant activity, resulting from the presence of some flavonoids in the extracts.

Contribution

First two authors contributed equally in this study.

Acknowledgements

The present study was supported by the Applied Basic Research Programs Foundation of Hunan, China (11K052), Hunan Higher foster industrialization projects (13cy026); The Technology R & D Program of Hunan Province, China (2013FJ6090).

References

- Altas S, Kizil G, Kizil M, Ketani A, Haris PI. Protective effect of Diyarbakir watermelon juice on carbon tetrachloride-induced toxicity in rats. *Food Chem Toxicol.* 2011; 49: 2433-38.
- Atawodi SE, Adekunle OO, Bala I. Anti-oxidant, organ protective and ameliorative properties of methanol extract of *anogeissus leiocarpus* stem bark against carbon tetrachloride-induced liver injury. *Int J Pharm Sci Res.* 2011; 2: 748-53.
- Basu S, Haldar N, Bhattacharya Sanji, Biswas S, Biswas M. Hepatoprotective activity of *Litchi chinensis* leaves against paracetamol-induced liver damage in rats. *Am-Eur J Sci Res.* 2012; 7: 77-81
- Balamurugan G, Muthusamy P. Observation of the hepatoprotective and antioxidant activities of *Trianthema*

- decandra* Linn. (Vallai sharunnai) roots on carbon tetrachloride-treated rats. Bangladesh J Pharmacol. 2008; 3: 83-89.
- Dong HQ, Feng ML, Liu FL, Huang JB. Inhibitory potential of trilobatin from *Lithocarpus polystachyus* Rehd. against α -glucosidase and α -amylase linked to type 2 diabetes. J Ethnopharmacol. 2012; 130: 261-66.
- Gilani AH, Jabeen Q, Ghayur MN, Janbaz KH and Akhta MS. Studies on the antihypertensive, antispasmodic, bronchodilator and hepatoprotective activities of the *Carum copticum* seed extract. J Ethnopharmacol. 2005; 98: 127-35.
- Halliwell B, Gutteridge JMC. Role of free radicals and catalytic metal ions in human disease: An overview. Method Enzymol. 1990; 186: 59-85.
- He NW, Yang XB, Jiao YD, Tian LM, Zhao Y. Characterisation of anti-oxidant and anti-proliferative acidic polysaccharides from Chinese wolfberry fruits. Food Chem. 2012; 133: 978-89.
- Hou SZ, Xu SJ, Jiang DX, Chen SX, Wang LL, Huang S, Lai XP. Effect of the flavonoid fraction of *Lithocarpus polystachyus* Rehd. on spontaneously hypertensive and normotensive rats. J Ethnopharmacol. 2012; 143: 441-47.
- Institute of Botany, Chinese Academy of Sciences. Iconographia Cormophytorum Sinicorum. Science Press, Beijing, 1972, p 434.
- Jain A, Soni M, Deb L, Jain A, Rout SP, Gupta VB, Krishna KL. Anti-oxidant and hepatoprotective activity of ethanolic and aqueous extracts of *Momordica dioica* Roxb. leaves. J Ethnopharmacol. 2008; 115: 61-66.
- Li SH, Wu XJ, Zeng JY, Zhang J, Zhou N. Studies on the flavonoids constituents of *Lithocarpus polystachyus*. Chinese Traditional and Herbal Drugs. 2010; 41: 1967-69.
- Li SH, Wu XJ, Yu JP, Li S, Han YG. Study on extraction techniques and content variety trends of total flavonoids in *Lithocarpus polystachyus* Rehd.. Food Science. 2008; 29: 139-41.
- Lin HM, Tseng HC, Wang CJ, Lin JJ, Lo CW, Chou FP. Hepatoprotective effects of *Solanum nigrum* Linn. extract against CCl₄-induced oxidative damage in rats. J Chemo-Biol Interac. 2008; 171: 283-93.
- Lu XS, Zhao Y, Sun YF, Yang S, Yang XB. Characterisation of polysaccharides from green tea of Huangshan Maofeng with anti-oxidant and hepatoprotective effects. Food Chem. 2013; 41: 3415-23.
- Nisar M, Qayum M, Cavar S, Shah MR, Zia-Ul-Haq M, Khan I, Ahmad KW, Qayum ZA. Chemical constituents and anti-oxidant activity of n-hexane extract of *Impatiens bicolor*. Chem Nat Comp. 2012; 48: 143-46.
- Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C. Anti-oxidant activity applying an improved ABTS radical cation decolorization assay. Free Radic Biol Med. 1999; 26: 1231-37.
- Recknagel RO. A new direction in the study of carbon tetrachloride hepatotoxicity. Life Sci. 1983; 33: 401-08.
- Singh N, Kamath V, Narasimhamurthy K, Rajini PS. Protective effects of potato peel extract against carbon tetrachloride-induced liver injury in rats. Environ Toxicol Pharmacol. 2008; 26: 241-46.
- Srinivasan R, Chandrasekar MJN, Nanjan MJ, Suresh B. Anti-oxidant activity of *Caesalpinia digyna* root. J Ethnopharmacol. 2007; 113: 284-91.
- Srivastava A, Shivanandappa T. Hepatoprotective effect of the root extract of *Decalepis hamiltonii* against carbon tetrachloride-induced oxidative stress in rats. Food Chem. 2010; 118: 411-17.
- Wang DY, Zhao Y, Jiao Y D, Yu LH, Yang S, Yang XB. Anti-oxidative and hepatoprotective effects of the polysaccharides from *Zizyphus jujube* cv. Shaanbeitanzao. Carbohydrate Polymers, 2012; 88: 1453-59.
- Weber LWD, Boll M, Stampfl A. Hepatotoxicity and mechanism of action of haloalkanes: Carbon tetrachloride as a toxicological model. Crit Rev Toxicol. 2003; 33: 105-36.
- Xu JY, Su YY, Cheng JS, Li SX, Liu RL, Li WX, Liu RL, Li WX, Xu GT, Li QN. Protective effects of fullerene on carbon tetrachloride-induced acute hepatotoxicity and nephrotoxicity in rats. Carbon 2010; 48: 1388-96.
- Zhou G, Chen YX, Liu S, Yao XC, Wang YW. *In vitro* and *in vivo* hepatoprotective and anti-oxidant activity of ethanolic extract from *Meconopsis integrifolia* (Maxim.) Franch. J Ethnopharmacol. 2013; 148: 664-70.
- Zhou DN, Ruan JL, Cai YL, Xiong ZM, Fu W, Wei AH. Anti-oxidant and hepatoprotective activity of ethanol extract of *Arachniodes exilis* (Hance) Ching. J Ethnopharmacol. 2010; 129: 232-37.

Author Info

Xianjin Wu (Principal contact)

e-mail: hhuxianjin@163.com