

CASE REPORTS

PENILE FRACTURE-REPORT OF TWO CASES AND REVIEW OF CURRENT LITERATURE

ASHRAF UDDIN MALLIK¹, MOSTAFIZUR RAHMAN², UTTAM KARMAKER³, BAIKALI FERDOUS⁴, HARBILASH HALDER⁵

Abstract

Penile fracture is an uncommon urological emergency, especially in Bangladesh. The other name is traumatic rupture of the tunica albugenia and corpora cavernosa in the erect penis. It occurs when an erect penis face to buckle under the pressure of a blunt sexual trauma. Patient gives the typical history of immediate detumescence, severe pain, swelling and eggplant deformity of the penile shaft due to penile injury. Immediate surgical exploration and repair of corpora Cavernosa with tunica albugenia is the most effective treatment modality. In normal cases diagnosis is made from history, physical examination alone. In some special cases ultrasonogram, radiological images, including retrograde urethrography or cavernosography are mandatory for proper diagnosis.

Key words: Cavernosography, Penile fracture, Tunica Albugenia rupture, Urethrography.

Bangladesh J. Urol. 2019; 22(1): 95-99

Introduction

Rupture of the tunica albuginea, which covers the penile shaft along with either one or both corpora cavernosa muscles of the penis is known as penile fracture. This fracture may be associated with injury to corpora spongiosum penis or urethra. This type of injury occurs when the erect penis is accidentally hits on the symphysis pubis or outside vagina after slipping out of the vaginal canal during coitus[1]. Partial or complete rupture of the urethra or injury to the dorsal nerves, veins, and arteries are sometimes involved[2]. It is a urological emergency that always deems attention[3]. Although penile fracture is easily recognized and therefore classified as a “first-look diagnosis,” this clinical entity is always embarrassing for patients as well as partners and goes unreported many times[4]. Eminently it is an entity of clinical diagnosis[5]. Therefore the management of a penile fracture should not include any further investigation rather than surgical exploration. The need for immediate

surgery is emphasized, in order to avoid erectile failure and curvature. Many conditions can simulate fracture penis as dorsal vein tears[6,7,8]. Authors reviewed the literature of penile fracture and reported two cases of 21 and 50-year-old men who presented with eggplant deformity of penile shaft and discoloration of the penile skin and swelling penis after cracking sound during forceful bending of penile shaft followed by severe pain at the time of fracture. The main aim of this study was to describe treatment option of 2 patients with fracture penis in our urology department and review of the literature.

Case 1

We report the case of a 21-year-old unmarried student who presented to the emergency department few hours following blunt injury to the penis during forceful bending of the erect penis. The patient reported a “tearing/popping” sensation, rapid detumescence, severe penile pain and no blood per urethra when he tried to bend the erect penis by himself. Following the injury, the patient was able to void without any voiding sensation. On physical examination the penis was found swollen, discoloration of the penile skin, scrotum, and part of the perineum, eggplant deformity of penile shaft (Figure 1). “Rolling sign” was present bilaterally, palpated at the right sites of tunica albuginea disruption about 1 cm from the base of the penis and the penis was not tender on examination. Clinically diagnosed with the penile fracture.

1. Professor, Department of Urology, KYAMCH, Enayetpur, Sirajgonj, Bangladesh.
2. Assistant Professor, Department of Urology, KYAMCH, Enayetpur, Sirajgonj, Bangladesh
3. Associate Professor, Department of Urology, DMCH, Dhaka, Bangladesh.
4. Register, Department of Urology, KYAMCH, Enayetpur, Sirajgonj, Bangladesh.
5. Consultant, Department of Anesthesiology, KYAMCH, Sirajganj, Bangladesh.

Correspondence : Dr. Ashraf Uddin Mallik, Prof. of Urology, KYAMCH, Enayetpur, Sirajgonj, Bangladesh, Phone: +8801928379099, e-mail: ashrafmallik143@gmail.com



Fig.-1: Eggplant deformity of penile shaft after penile fracture

Case 2

A 50-year-old married business male presented with a history of bending of his penis during vigorous coitus with his wife after taking an erotic stimulant drug at 12.30 night. During coitus he felt sudden thrust with the body of his wife followed by cracking sound, sudden detumescence, swelling of scrotum and deformity of the penis. On physical examination, the penis was swollen, discoloration of the penile skin, eggplant deformity of penis and "Rolling sign" was present. On the basis of his clinical presentation, a diagnosis of penile fracture was made. A penile catheter of 16F was inserted per urethra to serve as stent. Vital signs were normal. Pathological examinations like CBC, BT, CT, S.creatinine, RBS were done. Immediate surgical exploration and repair of rupture tunica albuginea and corpora cavernosa were done under spinal anesthesia. In both cases, a sub coronal circumscribes degloving incision was given. A tear in tunica albuginea and corpora cavernosa was found on the left side and 2 cm distal to the root of the penis and there was a blood clot. Repair of corpora cavernosa and tunica albugenia with sutures of vicryl 2-0 was performed after removing the clot (Figure 2). Post-operatively patient was prescribed antibiotics, analgesic and the urethral catheter was kept for 5 days, which was subsequently removed before discharge (Figure 3). Both of them were asked to abstain from masturbation or intercourse until complete healing occurred. At 3 months interval the optimal voiding function, erection function, and cosmetic result were achieved in one case and in another patient urethral stricture developed, which was repaired successfully after 4 months. Based on our experience, this management approach results an excellent preservation of both penile anatomy and function.



Fig.-2: Fracture of Tunica Albuginea and Corpora Caverosa



Fig.-3: Just after repair of penile fracture

Discussion

Penile fracture is a rare urologic emergency condition that affects all social strata. It has an incidence rate of 1 in 175,000 admissions and is commoner in countries where there is segregation of the sexes for social and religious reasons, where its etiologic factor is mostly from self manipulation[9,10].

Mechanism of fracture: During an erection, the penis is engorged with blood. If the penis is bent suddenly or forcefully while it's engorged, the trauma may rupture tunica albuginea, the lining of corpora cavernosa responsible for erections - resulting penile fracture. As the penis changes from a flaccid state to an erect state, the thick tunica albuginea becomes very thin. The

tunica albuginea is the fibrous covering of the penile corpora cavernosa and is directly involved in maintaining an erection. During erection it thins from 2 mm to 0.25 - 0.5 mm, stiffens and becomes less elastic and easily fracturable. Sudden direct trauma or abnormal bending in erect state causes tearing of the tunica albuginea, usually in transversal direction, rarely oblique or irregular[5].

Penile fractures have been classified as simple and compound[11]. Simple penile fractures are those with intact skin and urethra while those with urethral rupture are compound. One patient treated in our department had simple fractures another was compound fracture involving the urethra which was reported later having urethral stricture that was corrected surgically. Concomitant urethral injury occurs in about 10-58% of cases¹². Associated urethral rupture is seen more in coital fractures than those following manipulations that was found in our one case also[11]. The predisposing factor to concomitant urethral injury in penile fracture is not clear but it is postulated to be due to a more vigorous force applied during coitus compared with the prolonged but lesser force involved in masturbation[11,13]. Penile fracture may present with classic "eggplant deformity" of swollen penis along with ecchymosis confined to Buck's fascia[11]. The patient usually describe a cracking or popping sound during injury as tunica tears, followed by pain, rapid detumescence, discoloration and swelling of penile shaft[1,2,3]. Fracture typically occurs during vigorous sexual intercourse, when the erect penis slips out of the vagina and strikes the perineum or pubic bone, Other causes maybe masturbation with or without devices. It can happen from any type of blunt trauma affecting the tumescent shaft. Falling out of bed with an erection, extreme sexual activity, especially during coitus in which the female is on top, forceful correction of a congenital chordee and even tucking an erect penis into underwear. In the Middle East self-inflicted fractures predominate. Taqaandan also a cause of penile fracture. It comes from a Kurdish word meaning "to click," involves bending the top part of the erect penis while holding the lower part of the shaft in place, until a click is heard and felt. Penile fracture can usually be diagnosed based solely on history and physical examination findings. Sometimes in complicated cases, and to find out associated injury ultrasonogram, urethrogram, cavernosogram, MRI should be performed. Because of fear and embarrassment the patient present to health care professional sometimes

significantly in delay. But in our case both the victim gave detail history when we asked. False fracture penis has been reported in the literature, who presented with penile swelling and ecchymosis. In those case, they did not describe classic "snap-pop" or rapid detumescence "egg-plant" deformity of the shaft, whose features are associated with fracture penis. Physical examination may not be adequate for confirming diagnosis. Dorsal penile artery or vein injury during sexual intercourse, mimic penile fracture. Lacerations of the corpus cavernosa from gunshot and sporting injuries to the flaccid penis are not considered as penile fractures since they lack the fulcrum for snapping[12]. In western countries, up to 50% of penile fracture occurs during vigorous intercourse. Other causes include industrial accidents, masturbation, gunshot wound or other mechanical trauma that causing forceful breaking of an erect penis. In the Middle East, the injuries occur due to penile manipulation to achieve detumescence. Another causes are turning over in bed, a direct blow or forceful bending of the penis. In Nigeria reported causes of fracture are masturbation, stuttering priapism, vigorous sexual intercourse, turning in bed, forceful bending in the erect penis by the locally made bamboo bed[13]. The recent finding indicates that penile fracture appears to be in those population who use sexual exciting drugs before sexual intercourse. The case reported above was having coitus with wife using sexual stimulant drugs and another was masturbation. Treatment may be either conservative or surgical. The conservative management of penile fracture includes splinting cold compresses, and a combination of anti-inflammatory, analgesic medications and fibrinolytic. In our 2 cases, we did surgical exploration immediately after fracture and repaired. This concept has fallen into disfavor because of the high complication rates (29-53%) of nonoperative therapy[14].

Commonly complications encountered following conservative management as reported in literatures are penile angulation, painful erection, and coitus, A-V fistula, infected hematoma, abscess formation and impotence[3]. For these complications immediate surgical intervention is advocated as we did in our cases[15]. The goal of immediate surgical correction to the fracture a penis is restoration of the penis to its pre-injury state, prevent erectile dysfunction, maintain penile length and allow normal voiding[3]. It causes short-term hospital stay, patient satisfaction, safe from penile deformity development

and erectile dysfunction. Ideal management of penile fracture include identification of proper injury site, evacuation of hematoma, removal of corporal debridement, properly closure of tunica albuginea, corpora and ligation of any bleeding vessels[16]. Three types of incisions are advocated direct incision over

corpus cavernosum causes minimal dissection of the neurovascular bundles, less trauma, and even local anesthesia can be used but it does not allow complete repair. Some did repair with general anesthesia, but in our cases we did a repair with spinal anesthesia and achieved a good result. There was no impaired penile sensation or distal skin necrosis as described by others. That's why sub-coronal incision with spinal anesthesia appears to be satisfactory than other incisions. Treatment options for partial urethral tears include urethral catheterization, primary closure with non-absorbable suture, or suprapubic cystostomy tube. A penile fracture that may have devastating physiologic and psychologic consequences. However, with prompt diagnosis and expedient surgical management, outcomes remain excellent and complications are minimal[17,18,19].

Conclusion

The diagnosis of penile fracture depends on history and physical examination, but to find out injury to urethra imaging study is recommended. If there is Hematuria or urinary symptoms, it does not mean urethral injury. In that case retrograde urethrography is mandatory. Most of the time, penile fracture occurs on the lateral side of the proximal corpora but can occurs any portion of the penile shaft and can produce a variety of deformity. As penile fracture is a urological emergency, therefore, repair should be done promptly. It has been suggested that immediate surgical exploration can decrease complications and improve functional status. Conservative treatment may be given in properly selected patients. Studies with large number of patients and long term follow-up is necessary, to see the long term sequelae of fracture penis, is recommended.

References

1. *Greenberg's Text-Atlas of Emergency Medicine. Lippincott Williams & Wilkins. 22 November 2004.*

p. 318. ISBN 978-0-7817-4586-4. Retrieved 15 October 2012.

2. Haas CA, Brown SL, Spirnak JP "Penile fracture and testicular rupture". *World J Urol.* 1999; 17 (2): 101–6.
3. Mahapatra RS, Kundu AK, Pal DK. Penile Fracture: Our Experience in a Tertiary Care Hospital. *World J Mens Health.* 2015 ; 33 (2) : 95-102.
4. Bhoil R, Sood D. Signs, symptoms and treatment of penile fracture. *Emerg Nurse.* 2015; 9. 23 (6):16-7.
5. Swanson DE, Polackwich AS, Helfand BT et.al . Penile fracture: outcome of early surgical intervention. *Urology.* 2014; 84(5) : 1117-21 .
6. Perovic SV, Djinojic RP, Bumbasirevic MZ, Santucci RA, Djordjevic ML, Kourbatov D. Severe penile injuries: a problem of severity and reconstruction. *BJU Int.* 2009; 20.
7. Ghilan AM, Al-Asbahi WA, Ghafour MA, Alwan MA, Al-Khanbashi OM. Management of penile fractures. *Saudi Med J.* 2008; 29(10):1443-7.
8. Maruschke M, Lehr C, Hakenberg OW. Traumatic penile injuries—mechanisms and treatment. *Urol Int.* 2008; 81(3):367-9.
9. Rajih E, Alenizi A, El-Hakim A. Penile fracture with two epsilateral Corporal tears and delayed presentation : A case report. *Can Urol Assoc J.* 2015 ; 9(9-10).
10. Majzoub AA, Canguven O, Raidh TA. Alteration in the etiology of penile fracture in the Middle East and Central Asia regions in the last decade ; a literature review. *Urol Ann.* 2015 ; 7(3) : 284-8 .
11. Eke N. Fracture of the penis. *Br J Surg.* 2002; 89:555-565.
12. Koifman L, Cavalcanti AG, Manes CH et. al. Penile fracture. Experience in 56 cases. *Int. Braz J Urol* 2003; 29(1):35-39.
13. Al Ansari A, Talib RA, Shamsodini A et al. Which is guilty in self-induced penile fracture, marital Status, culture or geographic region ? A case series and literature reviews. *Int Impot Res.* 2013; 25(6):221-3.

14. Shittu OB, Kamara TB. Fracture of the penis diagnosis and management. *Afr J Med Sci* 2000; 29: 179-180.
15. Al-Reshaid RA, Madbouly K, Al-Jasser A. Penile abscess and necrotizing fasciitis secondary to neglected false penile fracture. *Urol Ann.* 2010;2:86–88.
16. Agarwal MM, Singh SK, Sharma DK, Ranjan P, Kumar S, Chandramohan V, et al. Fracture of the penis: a radiological or clinical diagnosis? A case series and literature review. *Can J Urol.* 2009; 16(2):4568-75.
17. Llarena IR, Villafruela MA, Azurmendi AJ, et al. Penile fracture with associated urethral rupture. *Arch Esp Urol.* 2006; 59:732-736.
18. Song W, Ko KJ, Shin SJ, Ryu DS. Penile abscess secondary to neglected penile fracture after intracavernosal vasoactive drug injection. *World J Mens Health.* 2012; 30(3):189-91.
19. Ateyah A, Mostafa T, Nasser TA, Shaeer O, Hadi AA, Al-Gabbar MA. Penile fracture: surgical repair and late effects on erectile function. *J Sex Med.* 2008 ; 5(6):1496-502.