

Original Article

Wealth Differentials in Prevalence of Self-Reported Diabetes Mellitus in Bangladeshi Adults

M Mostafa Zaman¹, Ferdous Hakim¹, Md Rijwan Bhuiyan², Md. Mujibur Rahman³, HAM Nazmul Ahsan³, Ridwanur Rahman⁴, Md. Abdul Jalil Chowdhury⁵

¹Research and Publication Unit, World Health Organization (WHO), Bangladesh, ²Research Officer, STEPS 2018, National Institute of Preventive and Social Medicine, Dhaka, ³Department of Medicine, Dhaka Medical College, ⁴Department of Medicine, Shaheed Suhrawardy Medical College, Dhaka, ⁵Department of Medicine, Bangabandhu Sheikh Mujib Medical University, Dhaka

Abstract:

Key words:
Diabetes, Wealth indices, Education, Bangladesh.

Background: We report relationship of wealth indices with self-reported diabetes data in Bangladeshi people. As a secondary objective we examine whether educational achievement can be used as a proxy of wealth indices.

Methods: We have analyzed data of a nationally representative STEPS survey (STEPwise Surveillance of NCD risk factors) done in 2010 in 8808 adults selected at their households using a multistage cluster sampling approach to describe association of socioeconomic status with diabetes mellitus. We created wealth indices using principal component analysis based on a 20-item household asset items; and then divided subjects in to quartiles.

Results: Prevalence of documented diabetes was 3.9% but it was more than 11 times (9.3%) in the fourth quartile (richest) compared to the first quartile (0.8%). This may either mean that the detection rate is higher among the higher socioeconomic group or there are some intervening factors accompanied by the economic achievements that favors occurrence or detection of diabetes in Bangladeshi adults.

Conclusion: Future studies need to identify the factor(s) that provides such a putative effect. Educational achievement also has shown similar trend. Therefore, education might be a proxy variable to determine socioeconomic status instead of a long list of household asset items.

(*Cardiovasc. j.* 2020; 13(1): 52-55)

Introduction:

There are lines of evidences that the prevalence of diabetes mellitus is rising in Bangladesh¹

as is happening in the South-East Asia Region² and elsewhere³. Diabetes is a major cause of death, disability and shorter lifespan.⁴ Therefore its control has been considered for noncommunicable diseases (NCD) monitoring framework of the World Health Organization⁵ and Sustainable Development Goals⁶. The occurrence of diabetes reported to have a socioeconomic

differential. There are studies that diabetes happens more in

poor⁷ while others report that it as a disease of the rich⁸. In this paper we report relationship of wealth indices with self-reported diabetes data in Bangladeshi people. As a secondary objective we examine whether educational achievement can be used as a proxy of wealth indices.

Methods:

This analysis is based on a nationally representative STEPS survey 2010 done by

Address for Correspondence: Dr. M Mostafa Zaman, World Health Organization, United House, Dhaka, Bangladesh. Email: zamanm@who.int

© 2020 authors; licensed and published by International Society of Cardiovascular Ultrasound, Bangladesh Chapter and Bangladesh Society of Geriatric Cardiology. This is an Open Access article distributed under the terms of the CC BY NC 4.0 (<https://creativecommons.org/licenses/by-nc/4.0>)

Bangladesh Society of Medicine using a multi-stage cluster design. It was a household-based survey for those aged 25 years or older ($n=9\,275$). However, in the current analysis we include those aged 25 to 69 years ($n=8\,808$) for comparability with other studies elsewhere as per STEPS protocol.⁹ Details of methods including ethical clearance have been reported elsewhere.¹⁰ Briefly, data on education and 20-item household assets were collected to define socioeconomic status. We did not measure blood glucose, and therefore defined diabetes by current medication history (insulin or oral hypoglycemic agents). In doubtful cases enumerators inquired about the names of medicines and checked the prescriptions or strips (or injection vials) of medicines if necessary.

We have done a complex survey design analysis of data, using *Epi Info* statistical package version 7.2.2, to obtain prevalence (with 95% confidence intervals) of diabetes in age and sex groups. Wealth indices were created using principle component analysis of household asset items,¹¹ and subjects were categorized into quartiles. Finally, prevalence estimates and their 95% confidence intervals were plotted in column charts according to wealth quartiles in men and women.

Results:

Self-reported prevalence of diabetes was 3.9% (95% confidence interval, 3.4% – 4.5%). Prevalence estimates were not statistically different between men and women (4.2% vs 3.6%). Diabetes prevalence increased with age consistently (data not shown). Prevalence was almost three times higher in urban areas (5.8%) compared to rural areas (2.0%) which was statistically significant. The urban people were more overweight and less active compared to rural people (data not shown).

There has been a clear gradient (in increasing order) of diabetes across the wealth quartiles (Figure 1, panel a). Prevalence in fourth quartile was more than 11 times (9.3%) than first quartile (0.8%). We observed similar but a little subtle trend for educational achievements (Figure 1, panel b).




Fig-1: Prevalence of self-reported diabetes according to wealth quartiles (a) and educational achievements (b) in men and women

Discussion:

This is the first study done in adults aged 25 years or older to report socioeconomic gradients of diabetes mellitus, although we used a case definition that limits the scope of discussion. Our case definition (medication for diabetes) might have excluded some people those who were diagnosed but were not taking medicines. Use of blood glucose data could yield a higher prevalence of diabetes.¹² The only national study using blood glucose was done for those aged 35 years and above, although the details of selection for ages beyond 45 years are not available.⁸ Therefore, a survey of national scale using blood glucose measurement for ages 18 years and older in line with Sustainable Development Goals is warranted. The use of household assets to determine socioeconomic status is a common practice where income cannot be easily determined. One advantage of using assets is their visibility in addition to an indication how money earned is spent to improve livelihood. However, collecting data on dozens of items is cumbersome.

Respondent reaction to such asset inquiry is frequently unwelcoming. We need to explore some other variable that would be time efficient to avoid fatigue from either side. One such variable, for Bangladeshi people, could be educational achievement. We have seen similar gradients of diabetes across educational categories. Our data on hypertension also have shown similar trend (not shown). Therefore, we suggest that educational achievement, at least for a medium term, could be used as a proxy of socioeconomic status ascertained out of a long list households asset items.

We acknowledge that our definition of diabetes might have favored inclusion of relatively rich people as numerator of current analysis. Keeping this in mind, we report here that economic achievement is linked to diabetes. Whereas in some developed countries it is the poor who suffer most.¹³ We need to find out which factor(s) is responsible for this opposite trend and spoiling the fruit of economic emancipation in our society.

Our data supports that a higher rate of diabetes in urban area is linked to overweight and insufficient physical activity.¹⁰ This might be due to some obesogenic diet and some other stressors for which we do not have data at hand.¹⁴ Urban people are generally more aware about health issues because of higher rates of education and media coverage. Therefore, increasing awareness in general is not the answer to this paradox. Specific points have to be identified for intervention at least to 'halt rise in diabetes'.⁵

Conclusion:

We would like to say that economic emancipation is warranted but factors intervening between this achievement and occurrence of diabetes must be explored before it is too late. Development initiatives and urban planning employed in Bangladesh needs a revisit to control diabetes. Similar finding of wealth indices and educational achievement to reflect socioeconomic status provides an indication that educational achievement can be used as a proxy variable at least for a medium term.

Declarations:

Competing interests: The authors declare that they have no competing interest. The views expressed

in this article are solely of the authors and they do not necessarily represent the views, decisions or policies of the institutions with which they are affiliated with. Funding: WHO, Bangladesh Country Office (WHO Reference: 2010/60773-1). Authors' contribution: MMZ conceived of the presented idea. MMZ and FH developed the theory and MRB performed the computations and referencing. MMZ, MMR, MRR and MAJC verified the analytical methods. All authors discussed the results and contributed to the final manuscript.

Conflict of Interest - None.

Acknowledgement:

The author gratefully acknowledges investigators of Bangladesh Society of Medicine and NCD Team of WHO Bangladesh for their contribution.

References:

1. Biswas T, Islam A, Rawal LB, Islam SMS. Increasing prevalence of diabetes in Bangladesh: a scoping review. *Public Health* 2016 Sep; 138:4–11. DOI: 10.1016/j.puhe.2016.03.025
2. Lim SS, Vos T, Flaxman AD, Danaei G, Shibuya K, Adair-Rohani H, et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012 Dec 15;380(9859):2224–60. DOI: 10.1016/S0140-6736(12)61766
3. Guariguata L, Nolan T, Beagley J, Linnenkamp U, Jacqmain O. IDF Diabetes Atlas 6th edition [Internet]. Brussels: International Diabetes Federation; 2013 [cited 2019 Jan 2] p. 15–7. Report No.: 930229-85–3. Available from: <https://www.idf.org/elibrary/epidemiology-research/diabetes-atlas/19-atlas-6th-edition.html>
4. Bhutani J, Bhutani S. Worldwide burden of diabetes. *Indian J Endocrinol Metab* 2014; 18(6) :868–870. DOI: 10.4103/2230-8210.141388
5. WHO. Noncommunicable Diseases Global Monitoring Framework: Indicator Definitions and Specifications [Internet]. Geneva: World Health Organization; 2013 p. 3. Available from: https://www.who.int/nmh/ncdtools/indicators/GMF_Indicator_Definitions_Version_NOV2014.pdf
6. UN. The Sustainable Development Goals Report [Internet]. Geneva: United Nations Statistics Division; 2017. Report No.: 978-92-1-101368–9. Available from: <https://sdgactioncampaign.org/wpcontent/uploads/2017/07/TheSustainableDevelopmentGoalsReport2017.pdf>
7. Hwang J, Shon C. Relationship between socioeconomic status and type 2 diabetes: results from Korea National Health and Nutrition Examination Survey (KNHANES) 2010-2012. *BMJ Open* 2014 Aug 19;4(8):e005710. DOI: 10.1136/bmjopen-2014-005710

8. Akter S, Rahman MM, Abe SK, Sultana P. Prevalence of diabetes and prediabetes and their risk factors among Bangladeshi adults: a nationwide survey. *Bull World Health Organ* 2014 Mar 1;92(3):204-213A. DOI: 10.2471/BLT.13.128371
9. WHO. NCDs | STEPS conceptual framework [Internet]. WHO. [cited 2019 Jan 2]. Available from: <http://www.who.int/ncds/surveillance/steps/framework/en/>
10. Zaman MM, Rahman MM, Rahman MR, Bhuiyan MR, Karim MN, Chowdhury MAJ. Prevalence of risk factors for non-communicable diseases in Bangladesh: Results from STEPS survey 2010. *Indian J Public Health* 2016 Mar;60(1):17–25. DOI: 10.4103/0019-557X.177290
11. Rutstein S, Johnson K. The DHS Wealth Index. DHS Comparative Reports No. 6 [Internet]. Calverton: Demography and Health Survey: US AID; Available from: <https://dhsprogram.com/pubs/pdf/cr6/cr6.pdf>
12. Zaman MM, Choudhury SR, Ahmed J, Talukder MH, Rahman AHMS. Blood glucose and cholesterol levels in adult population of Bangladesh: Results from STEPS 2006 survey. *Indian Heart J* 2016 Feb; 68(1):52–56. DOI: 10.1016/j.ihj.2015.06.031
13. Black SA. Diabetes, Diversity, and Disparity: What Do We Do With the Evidence? *Am J Public Health*. 2002 Apr;92(4):543–8. PMID: 11919048
14. Kann PH, Münzel M, Hadji P, Daniel H, Flache S, Nyarango P, et al. Alterations of cortisol homeostasis may link changes of the sociocultural environment to an increased diabetes and metabolic risk in developing countries: a prospective diagnostic study performed in cooperation with the Ovahimba people of the Kunene region/ northwestern Namibia. *J Clin Endocrinol Metab* 2015 Mar;100(3):E482-486. DOI: 10.1210/jc.2014-2625.