

Faulty Breast Feeding Practice : A Risk Factor in Malnourished Children

Sukhendu Shekhar Sen^{1*}
Jhulan Das Sharma²
Dhananjay Das³
Shahed Iqbal³
Md Badruddoza³

¹Department of Paediatrics
Chittagong Medical College, Chittagong, Bangladesh.

²Department of Paediatrics
Southern Medical College, Chittagong, Bangladesh.

³Department of Paediatrics
Chattagram Maa Shishu-O- General Hospital
Chittagong, Bangladesh.

*Correspondence to:

Dr. Sukhendu Shekhar Sen
Assistant Professor
Department of Paediatrics
Chittagong Medical College, Chittagong, Bangladesh
Mobile : +88 01713207027
E-mail: sukhendusen@gmail.com

www.banglajol.info/index.php/CMOSHMCJ

Abstract

Background: The predominant form of malnutrition is commonly called protein-calorie malnutrition. Protein Energy Malnutrition (PEM) is still a major health problem in children of developing countries including Bangladesh. The causes of malnutrition are multifactorial including nutritional factors, socioeconomic factors, health status of the mothers and repeated infections in children. **Objective:** To explore the information regarding the breast feeding practices of children suffering from Protein-Energy Malnutrition. **Methods:** This case control study was conducted in Chittagong Medical College Hospital from November 2006 to April 2007. A total of 65 controls and 65 cases were selected consecutively for the purpose of the study. Their mothers were interviewed with help of structured questionnaire containing all the variables of interest to attain the study objectives. The test statistics used to analyze the data were descriptive statistics and Chi-square (χ^2) or Fisher's Exact Probability Test. **Results:** A significantly higher frequency of cases (67.7%) were given pre-lacteal feed, predominantly honey and sugar-water compared to control group (41.5%) ($p = 0.008$). Nearly 100% of controls were given colostrums compared to 75% of the cases. About one-third (31.3%) of the cases was exclusively breast-fed in comparison to 58.5% of the control group ($p = 0.003$). Over one-quarter (27.7%) of the control were breast-fed upto 6 months of age, as opposed to only 1.5% cases ($p < 0.001$). Duration of predominant breast feeding for more than 6 months of age was also significantly higher in control group than that in case group ($p = 0.001$). Breast milk substitutes demonstrate their significant presence in cases (38.5%) than that in controls (9%) ($p = 0.003$). **Conclusion:** The study showed that rejection of colostrums, practice of prelacteal feeding, delayed initiation of breast feeding, early cessation of exclusive breast feeding and use of formula milk all were significantly higher in the malnourished group of children than those in their normal counterpart.

Key words: Protein energy malnutrition; Breastfeeding; Risk factors.

INTRODUCTION

The term "Protein Energy Malnutrition" (PEM) is used to cover a spectrum of clinical pictures, ranging from frank kwashiorkor to severe marasmus. The description of kwashiorkor given by Williams in her first published article (1933) is as vivid and accurate today as it was more than 50 years ago, although she did not introduce the name kwashiorkor until her second paper in 1935. The earliest account of the syndrome was published by Hinazosa in Mexico in 1865¹.

The predominant form of malnutrition is commonly called Protein-Calorie Malnutrition (PEM) the most serious and widespread nutritional disorder known to medical and nutritional science. To a varying extent, but common to all these forms of protein-calorie malnutrition, are retarded growth and disorder tissue repair (Protein deficiency), and energy storage (Calorie deficiency). Marasmus is total inanition in the child with severe and continued restriction of both calories and protein as well as of all other nutrients. In kwashiorkor dietary protein is of poor quality and restricted in amount; calories are often restricted but may even be in excess of requirements².

Magnitude of the problem:

Malnutrition remains one of the most common causes of morbidity and mortality among children throughout the world³. Some 49% of the 10 million deaths among the under-five children each year in the developing world are associated with malnutrition. Those who survive are at risk of disability, stunted mental and physical growth and as a result retarded national socio-economic development⁴. It is estimated that, 26.7% of the under-five children in the developing countries are affected by protein energy malnutrition and more than two-thirds (70%) of those children live in Asia (of which 61% are in south central Asia)⁵. Malnutrition is a major public health problem in Bangladesh. The Child Nutrition Survey of Bangladesh 2000 showed that about 2.4% of children were severely malnourished while about 34.7% were moderately malnourished⁶.

Multifactorial cause of malnutrition:

Whatever the form of presentation of malnutrition (marasmus or kwashiorkor), the root problem is not necessarily a shortage of food. The causes of malnutrition are multifactorial, including nutritional factors, socioeconomic factors, health status of the mothers and repeated infections in children⁷.

Infection and malnutrition creates a vicious cycle. Diarrhoea, measles, whooping cough, tuberculosis and infestation with helminthes compromise the nutrition of the child. Poor immunization status resulting in the persistence of the vicious cycle of infection and malnutrition¹.

Faulty feeding practice and malnutrition:

Faulty Breast feeding practice, including rejection of colostrums is among the most important causes of malnutrition.

Good feeding practice is crucial for under 2 child survival. Exclusive breast feeding is most desirable for 0-6 months age group children. Weaning should be started at 6 months of age. During 6-23 months age breastfeeding and supplementary food (Khichuri, egg, fruit, juices, vegetables, mashed meat, fish successively) are normal feeding. Proper preparation is also important here⁸.

Current breast feeding recommendations⁹

1. Babies should be exclusively breast feed for atleast 6 months.
2. Babies should receive no infant formula or animal milk or drinks or foods other than breast milk until 6 months.
3. No limit should be placed on the number or length of breast feeds.
4. Give the baby no bottles, artificial feeds or pacifiers / dummies.
5. Babies should continue breastfeeding for up to 2 yrs with increasing amounts of complementary foods and cup feed liquids.

OBJECTIVES

General: To explore the information regarding the breast feeding practices of children suffering from Protein-Energy Malnutrition.

Specific: To identify the faulty breast feeding practices in malnourished children in a hospital set up.

MATERIALS & METHOD

A case control design was adopted to conduct the study.

Place and duration of study:

The study was conducted in the "Nutrition Block" of Department of Paediatrics, Chittagong Medical College Hospital during the period from November 2006 to April 2007.

Study population:

Study population was divided into cases and controls.

Case: Moderate to severely wasted children with or without bipedal nutritional oedema and age ranging from 6 – 59 months admitted in the "Nutrition Block" of Department of Paediatrics, Chittagong Medical College Hospital were selected as case. Oedema due to renal or liver cause were excluded from the study.

Control: Children of same age range as cases with mild or no wasting without bipedal nutritional oedema admitted in the same hospital for diseases other than malnutrition or attended at OPD or EPI centre were taken as controls.

Sample size: The sample size was determined using the following formula.

$$n = (Z^2 \times p \times q) / d^2, \text{ where } Z = \text{Standard normal deviate} = 1.96$$

$$p = \text{Prevalence of the disease or event of interest} = 0.15$$

(as the prevalence of moderate and severe wasting in under five children is 15%)

$$q = (1-p) = 0.85 \text{ and}$$

$$d = \text{Desired accuracy} = 0.05$$

Putting the values in the equation we got the required sample size,

$$n = 1.96^2 \times 0.15 \times (1-0.15) / (0.05)^2 = 196.$$

Due to limitation of time, purposive and convenient sampling was done. A total of 65 controls and 65 cases were selected consecutively for the purpose of the study. From the statistical point of view it is significant.

Variables studied:

The following variables were studied to address the study objectives:

1. Socio-demographic characteristics:

- Age of child.
- Sex of child.

2. Feeding history:

- Prelacteal meal
- Ever breast milk given
- Initiation of breast feeding
- Colostrums intake
- Exclusive breast feeding
- Duration of exclusive breast feeding
- Predominant breast feeding
- Duration of predominant breast feeding
- Breast milk substitution within 6 months of age

Classification of malnutrition:

- WHO classification

Data collection:

Taking written consent from the parents/attendants of each of the subjects included in the study, data were collected using a structured questionnaire (research instrument) containing all the variables of interest.

Data processing:

Collected data were checked and edited first. They were then processed with the help of software Statistical Package for Social Sciences (SPSS) version 11.5 and analysed.

Statistical analysis:

Statistical analyses were done with help of descriptive statistics and Chi-square (χ^2) or Fisher's Exact Probability Test. Data presented on categorical scale were compared between case and control group using Chi-square (χ^2) or Fisher's Exact Probability Test. Level of significance for all analyses was set at 0.05 and p-value < 0.05 was considered significant.

RESULTS

Table 1 : Distribution of age between cases and controls.

Age(months)	Group	
	Case (n = 65)	Control (n = 65)
6 - 12	27(41.5)	34(52.3)
13 – 18	13(20.0)	11(16.9)
19 – 24	5(7.7)	6(9.2)
>24	20(30.8)	14(21.6)




Figure 1 : Distribution of cases and controls by sex.

Table 2 : Comparison of prelacteal meal between cases and controls.

Prelacteal meal	Group		Odds Ratio (95% CI of OR)	p-value
	Case (n = 65)	Control(n = 65)		
None	21(32.3)	38(58.5)	2.92(1.60 - 5.51)	0.008
Prelacteal meal	44(67.7)	27(41.5)		

Table 3 : Comparison of initiation of breast feeding between groups.

Initiation of breast feeding	Group			p-value
	Case (n = 65)	Control (n = 65)	Odds Ratio (95% CI of OR)	
Ever breast milk given				
Yes	61(93.8)	65(100.0)		0.119
No	4(6.2)	00		
Initiation of breast feeding				
Within 24 hour of birth	45(73.5)	64(98.5)		
After 24 hours of birth	16(26.5)	1(1.5)	17.22(3.8-108.5)	< 0.001




Figure 2 : Distribution of cases and controls by colostrum feeding.

Table 4 : Comparison of exclusive breast feeding between groups.


Breast feeding	Group			p-value
	Case (n = 65)	Control (n = 65)	Odds Ratio (95% CI of OR)	
Exclusive breast feeding				
Yes	20(31.3)	38(58.5)		0.003
No	45(68.7)	27(41.5)		
Duration of exclusive breast feeding (months)				
< 6	11(55.0)	7(18.4)	5.57(2.79 - 11.20)	< 0.001
≥ 6	9(45.0)	38(81.6)		

Table 5 : Comparison of predominant & partial breast feeding between group.

Breast feeding	Group		p-value
	Case (n = 65)	Control (n = 65)	
Duration of predominant breast feeding			
< 6 months	20(30.8)	4(6.2)	
up to 6 months	4(6.2)	6(9.2)	
> 6 months	12(18.5)	12(18.5)	0.001
Partial breast feeding#			
Yes	21(34.4)	9(13.8)	0.003
No	44(65.6)	56(86.2)	

Table 6 : Comparison of breast milk substitute between groups.

Breast milk supplements within 6 month of age*	Group		p-value
	Case (n = 25)	Control (n = 9)	
Formula milk	19(76.0)	9(100.0)	< 0.001
Cow's milk	6(24.0)	0(0.0)	

**Figure 3 :** WHO classification of malnutrition.

DISCUSSION

Protein energy malnutrition is still a major health problem in children of developing countries including Bangladesh. The findings of the present study aimed at identifying the risk factors for faulty breast feeding practice, therefore, need to be compared and contrasted with the studies conducted to the same end at home and abroad.

As demographic characteristics of the subjects are considered, the mean age of the cases (19.7 ± 13.0 months) was found to be significantly higher than that of controls (13.5 ± 7.8 months) ($p < 0.01$), although the sex distribution between groups was almost homogeneous. Amin et al. found majority (88%) of the malnourished children below 3 years of age. However, 30% of the malnourished children in our study were above 2 years of age. Mollah et al. found the mean age 12.3 months^{10,11}.

A significantly higher frequency of cases (67.7%) were given pre-lacteal feed predominantly honey and sugar water compared to control group (41.5%) ($p = 0.008$). The rest 58.5% of the control group were not given any pre-lacteal feed. The widely practiced pre-lacteal feeding could be explained by the ritualistic feeding practices and lack of confidence of mothers about sufficiency of breast milk produced during early lactation¹².

However, no significant difference was observed between cases and control group with respect to breast milk ever given ($p = 0.119$). But in terms of initiation of breast feeding 80% of the mothers of control group initiated it within half an hour of birth of their babies compared to 44.3% of mothers of the case group ($p < 0.001$). Early initiation of breast feeding preferably within half an hour of birth help to develop strong suckling reflex and successful lactation¹³.

Nearly 100% of controls were given colostrum compared to 75% of the cases. The difference between cases and controls was statistically significant ($p < 0.001$). Faulty Breast feeding practice including rejection of colostrums is among the most important causes of malnutrition. Haque et al. showed that colostrums feeding was significantly low ($p < 0.01$) and pre-lacteal feeding was significantly high ($p < 0.01$) among the cases in comparison to the controls¹⁴.

About one-third (31.3%) of the cases was exclusively breast-fed in comparison to 58.5% of the control group ($p = 0.003$). Over one-quarter (27.7%) of the controls was breast fed up to 6 months of age, as opposed to only 1.5% cases ($p < 0.001$) which is consistent with the study of Mollah et al. Early cessation of Breast feeding is a major cause of severe protein-energy malnutrition among the under two children. This was shown in a recent study in Tanzania^{11,15}. Duration of predominant breast feeding for up to 6 months of age was also significantly higher in control group than that of case group ($p = 0.001$). Partial breast feeding demonstrates their significant presence in cases (38.5%) than that in controls (15.4%). Supplementation of breast milk with cow's milk and formula milk within 6 months of age was much higher in the case group than that in the control group ($p < 0.001$).

WHO classification of malnutrition shows that majority (85%) of the cases had severe malnutrition and rest 15% moderate malnutrition, while Welcome classification demonstrates that more than one-third (37%) of the cases had marasmas, 18% had kwashiorkor and 45% had marasmic-kwashiorkor. In another study 50% of the malnourished children had marasmus, 16% had kwashiorkor and 34% marasmic kwashiorkor¹⁶.

