



Study of Fine Needle Aspiration Cytology of Lymph Node Lesions of Head and Neck Region



Nazmun Nahar¹, Mohammed Iqbal², Kazi Md. Shahidur Rahman³, Rawshon Ara⁴,
Mst. Rommana Akter⁵, Raju Mollick⁶, Nishat Sultana⁷

¹Professor (cc), Department of Pathology, Medical College for Women & Hospital, Uttara, Dhaka, Bangladesh; ²Professor, Department of Anatomy, Monno Medical College, Manikgonj, Bangladesh; ³Associate Professor, Department of Pathology, Monno Medical College, Manikgonj, Bangladesh; ⁴Assistant Professor, Department of Anatomy, Monno Medical College, Manikgonj, Bangladesh; ⁵Assistant Professor, Department of Dermatology, Monowara Sikder Medical College, Shariatpur, Bangladesh; ⁶Assistant Professor, Department of Orthopaedic, Monowara Sikder Medical College, Shariatpur, Bangladesh; ⁷Assistant Professor, Department of Anatomy, Monowara Sikder Medical College, Shariatpur, Bangladesh;

Abstract

Background: Lymphadenopathy in the head and neck region is a common clinical issue with diverse causes, ranging from benign conditions to malignancies. Fine Needle Aspiration Cytology (FNAC) has become an essential diagnostic tool due to its minimally invasive nature, cost-effectiveness, and ability to provide rapid, accurate diagnoses. FNAC is particularly valuable for distinguishing between benign and malignant lymph node lesions and identifying specific infections like tuberculosis. **Objective:** This study focused on analyzing the frequency and types of lymph node lesions in the head and neck region using FNAC. **Methodology:** This prospective study, conducted from January 2022 to January 2023 in the Department of Pathology at the Medical College for Women & Hospital, included a total of 53 patients. The study focused on patients presenting with head and neck swelling. Data were collected using a pre-designed data collection form and analysed with SPSS software, Windows version 25. **Results:** Chronic nonspecific lymphadenitis was the most frequently observed pathological condition of the lymph nodes, comprising 50.9% of cases. Granulomatous lymphadenitis, including both tuberculous and non-tuberculous forms, was identified in 33.9% of cases, with the tuberculous form being more common. Acute suppurative lymphadenitis accounted for 13.2% of cases, while metastatic lesions, squamous cell carcinoma was the least common, each representing 1.9% of the total cases. The most common age group affected was 11-20 years (32%), and females were more frequently affected (71.7%). **Conclusion:** This study reveals that chronic nonspecific lymphadenitis is the most frequently encountered lymph node condition, followed by granulomatous lymphadenitis, particularly of tuberculous origin, acute suppurative lymphadenitis, and metastatic involvement (either squamous cell carcinoma or adenocarcinoma). Frequently female were affected and most common age group in between 11-20 years. FNAC is recommended as a safe, reliable minimally invasive method for diagnosing head and neck lesions. It serves as a complementary diagnostic procedure to histopathological examination in the diagnosis and management of palpable head and neck lesions. [Journal of National Institute of Neurosciences Bangladesh, July 2025;11(2):165-169]

Keywords: FNAC; Lymph Node; Head & Neck Region

Introduction

Lymph nodes play a crucial role in the human immune system. Lymphadenopathy arises when a lymph node becomes enlarged, increased in number, or exhibits atypical consistency^{1,2}. The causes of lymphadenopathy vary from benign reactive lymphoid hyperplasia to malignant conditions. While several conditions can cause

lymph node enlargement, the majority (90%) are due to benign causes, including reactive hyperplasia (60%) and infectious or inflammatory lymphadenitis (30%)^{2,3}. Infectious or inflammatory lymphadenitis encompasses conditions such as Kikuchi-Fujimoto disease (KFD), tuberculosis, sarcoidosis, infectious mononucleosis, toxoplasmosis, HIV infection, cat-scratch disease, and

Correspondence: Prof. Dr. Nazmun Nahar, Professor(cc), Department of Pathology, Medical College for Women & Hospital, Uttara, Dhaka; Email: nahar085081@gmail.com; Cell no +8801914140697;

ORCID: <https://orcid.org/0000-0002-6225-145X>

©Authors 2025. CC-BY-NC

drug reactions (e.g., to phenytoin). Malignant lymphadenopathies make up only 10.0% of cases, with primary lymphomas accounting for 3.0% (including diffuse large B-cell lymphoma, follicular lymphoma, mantle cell lymphoma, peripheral T-cell lymphoma, and Hodgkin's lymphoma) and metastatic carcinomas constituting 7.0% (including metastatic papillary thyroid carcinoma, adenocarcinomas from the lung or breast, and poorly differentiated squamous cell or small cell carcinomas)³.

Fine Needle Aspiration Cytology (FNAC) is widely used as a primary diagnostic method for evaluating enlarged lymph nodes and ruling out involvement of other organs, such as the salivary glands, head, neck, or other subcutaneous masses. This minimally invasive technique offers quick diagnosis and treatment with minimal complications, such as hemorrhage, nerve damage, and vasovagal reactions, particularly in head and neck procedures^{4,5,6}. FNAC is especially beneficial in developing countries where surgical biopsy costs may be prohibitive. It aids surgeons in selecting, guiding, and adjusting surgical plans or clinical management, including antibiotic therapy or neoadjuvant chemotherapy. This study aims to assess the use of FNAC for lymph node lesions in the head and neck region.

Methodology

The prospective study was conducted from January 2022 to January 2023 in the department of pathology of the medical college for women & hospital. Total 53 patients were included in this study. All patients presenting with head and neck swelling were included in the study, while those with masses located elsewhere were excluded. FNAC was performed under strict aseptic conditions using a 10 ml syringe with a 23 or 24

gauge needle, applying both aspiration and non-aspiration techniques. The collected material was placed on glass slides, which were then prepared and stained with Papanicolaou (PAP) stains. Data were gathered using a pre-designed collection form and analyzed with SPSS software, Windows version 25.

Results

In table 1 showed chronic nonspecific lymphadenitis is the most common pathological condition of the lymph nodes, accounting for 50.9% of cases. Granulomatous lymphadenitis, including both tuberculous and non-tuberculous forms, was observed in 33.9% of cases, with the tuberculous form being more prevalent. Acute suppurative lymphadenitis was present in 13.2% of cases, while metastatic lesions, squamous cell carcinoma was the least common, each making up 1.9% of the total cases.

Table 1: Place of Birth of Respondents

Variables	Frequency	Percent
Residence Wise		
• Urban	171	68.4
• Rural	79	31.6
Delivery Wise		
• Home	140	56.0
• Hospital	110	44.0

In Table 2 it is illustrated the analysis of 53 cases of lymph node lesions from the head and neck region indicates specific trends related to age and sex. The age group 11 to 20 years displays the highest frequency of lymphadenitis cases (32.0%), with chronic nonspecific lymphadenitis (13 cases) being the most common lesion in this category. Among patients aged 21 to 30 years, granulomatous lymphadenitis, including tuberculous

Table 2: Frequencies of Various Pathological Conditions of Lymph Nodes in Relation to Age and Gender (N=53)

Age of the Patient (in years)	Acute Suppurative Lymphadenitis	Chronic Nonspecific Lymphadenitis	Granulomatous Lymphadenitis (tuberculous)	Granulomatous Lymphadenitis	Metastatic (Squamous cell carcinoma)	Total	%
1 to 10	1	5	1	0	0	7	13.2
11 to 20	1	13	3	0	0	17	32.0
21 to 30	3	3	5	4	0	15	28.3
31 to 40	2	3	4	0	0	9	17.0
41 to 50	0	2	0	1	0	3	5.7
51 and above	0	1	0	0	1	2	3.8
Gender							
Male	0	12	2	0	1	15	28.3
Female	7	15	11	5	0	38	71.7

types, is notably prevalent, comprising 28.3% of cases. Acute suppurative lymphadenitis is mainly seen in patients aged 21 to 30 years. Metastatic lymphadenopathy, particularly in patients aged 51 and above, is rare but present (1.9% of cases), indicating a greater risk of malignancy with advancing age. Out of 53 patients, 15 (28.3%) were males and 38 (71.7%) were females. Chronic Nonspecific Lymphadenitis were highest frequency in female (28.3%) and male (22.6%)

Discussion

Head and neck lesions are frequently encountered in clinical practice. In this study, lymph nodes were the most common sites for head and neck masses, followed by the thyroid, other soft tissues, and salivary glands. These findings align with other studies, which also report lymph nodes as the primary site for FNAC in the head and neck region^{7,8,9}.

This study shows chronic nonspecific lymphadenitis emerges as the most prevalent condition, accounting for 50.9% of the cases, suggesting that it is the predominant cause of lymph node enlargement. Granulomatous lymphadenitis, particularly the tuberculous variant, is the second most common condition, comprising 24.5% of the cases. The presence of non-tuberculous granulomatous lymphadenitis, albeit less frequent at 9.4%, indicates the necessity to consider a broad differential diagnosis for granulomatous lymphadenitis, including sarcoidosis and fungal infections. Acute suppurative lymphadenitis, observed in 13.2% of the cases, reflects acute bacterial infections, which are a significant but less common cause of lymph node pathology compared to chronic conditions. The low frequency of metastatic involvement (1.9%) highlights the importance of lymph node examination in the staging and diagnosis of malignancies. This study points towards a predominance of benign, inflammatory conditions, with a notable proportion of cases attributable to tuberculosis. The relatively low incidence of metastatic disease suggests that while malignancy is a critical consideration, it is less commonly encountered in this clinical context. These findings emphasize the need for a thorough and differential diagnostic approach in the evaluation of lymphadenopathy. This findings consistent with other study Padia and Dhokiya⁵.

In a similar study, Pathak et al.⁷ reported that among 128 cases of lymph node lesions, reactive lymphadenitis was the most frequent pathological finding, followed by metastatic carcinomas,

granulomatous lymphadenitis, lymphomas, and abscesses. Other studies have also found reactive lymphadenitis to be the most common diagnosis for lymph node lesions, consistent with our findings¹⁰⁻¹².

In contrast, some studies have identified tuberculous lymphadenitis as the most common finding among lymph nodes^{1,2,5,6}. Among 66 cases available for histopathological examination, non-Hodgkin lymphoma (27.3%) was the most frequent pathological finding, followed by granulomatous lymphadenitis (25.8%), metastatic carcinoma (19.7%), reactive lymphadenitis (10.6%), Hodgkin lymphoma (7.6%), Kikuchi-Fujimoto disease (4.5%), and abscesses (4.5%). Another study by Shekha et al¹³ reported that reactive lymphadenitis was the most common pathology, accounting for 16.0% of cases, followed by granulomatous lesions (tuberculous) at 15.0%, and malignant lesions at 14.5%.

In a study by Sreedevi et al¹⁴ in Andhra Pradesh, lymph nodes accounted for 50.0% of head and neck lesions out of 304 cases, with reactive lymphadenitis being the most common lesion. Thyroid lesions were the next most frequent, with colloid goiter being the predominant diagnosis. Kaur et al¹⁵ reported that lymph node lesions comprised 52.89% of all head and neck lesions, with reactive lymphadenitis being the most common (33 cases, 25.78%), followed by tuberculous lymphadenitis (32 cases, 25%) and granulomatous lesions (24 cases, 18.75%). Other lesions included abscesses (10 cases, 7.81%) and benign cystic lesions (three cases, 2.34%). Among malignant lesions, metastases from squamous cell carcinoma were most prevalent (18 cases, 14.06%), followed by non-Hodgkin lymphoma (three cases, 2.34%) and one case of Hodgkin lymphoma (0.78%). Four cases (3.12%) were deemed inadequate due to hemorrhagic aspirate. In present metastatic squamous cell carcinoma was common than other malignant tumor (1.9%) which is similar with that study Goswami et al¹⁶ found tuberculous lymphadenitis to be the most common lymph node lesion (42.12%), while it was the second most common in our study (27.4%) and in Patel et al¹⁷ study (25.0%).

In the present study, lymphadenopathy was more common in females (71.7%), similar to findings by Kumar et al¹⁸ (64%). Mali et al¹⁹ reported that the most common age group for lymphadenopathy was 11 to 20 years (28.8%), which aligns with our findings, where the same age group accounted for 32.0%.

Conclusion

In conclusion, the relationship between educational This study indicates that the frequency distribution of various pathological conditions of lymph nodes shows chronic nonspecific lymphadenitis as the most common, followed by granulomatous lymphadenitis, particularly of tuberculous origin, granulomatous lymphadenitis, acute suppurative lymphadenitis, and metastatic involvement (squamous cell carcinoma). Frequently female were affected and most common age group in between 11-20 years. FNAC is recommended as a safe, reliable, quick, cost-effective, and minimally invasive technique for diagnosing head and neck lesions. It acts as a complementary diagnostic tool to histopathological examination in the diagnosis and management of palpable head and neck lesions.

Acknowledgements

None

Conflict of interest: All authors declare that they have no competing interest.

Financial Disclosure

This research project was not funded by any organization.

Contribution to authors: Conception and design: Nahar N, Iqbal M, Rahman KMS; Acquisition, analysis, and interpretation of data: Rahman KMS, Ara R, Akter MR, Mollick R; Data collection, consultation, Nahar N, Manuscript drafting and revising it critically: Ara R, Akter MR, Mollick R, Sultana N; All authors read and approved the final manuscript.

Data Availability

Any inquiries regarding supporting data availability of this study should be directed to the corresponding author and are available from the corresponding author on reasonable request.

Ethics Approval and Consent to Participate

Ethical approval for the study was obtained from the Institutional Review Board. As this was a prospective study the written informed consent was obtained from all study participants. All methods were performed in accordance with the relevant guidelines and regulations.

How to cite this article: Nahar N, Iqbal M, Rahman KMS, Ara R, Akter MR, Mollick R, Sultana N. Study of fine needle aspiration cytology of lymph node lesions of Head and Neck Region. *J Natl Inst Neurosci Bangladesh*, 2025;11(2):165-169

Copyright: © Nahar et al. 2025. Published by Journal of National Institute of Neurosciences Bangladesh. This is an open access article and is licensed under the Creative Commons Attribution Non-Commercial 4.0 International License (CC BY-NC 4.0). This license permits others to distribute, remix, adapt and reproduce or changes in any medium or format as long as it will give appropriate credit to the original author(s) with the proper citation of the original work as well as the source and this is used for noncommercial purposes only. To view a copy of this license, please See: <https://creativecommons.org/licenses/by-nc/4.0/>

ORCID:

Nazmun Nahar: <https://orcid.org/0000-0002-6225-145X>
 Mohammed Iqbal: <https://orcid.org/0000-0002-2346-361X>
 Kazi Md. Shahidur Rahman: <https://orcid.org/0000-0002-6493-334X>
 Rawshon Ara: <https://orcid.org/0009-0001-0581-640X>
 Mst. Rommana Akter: <https://orcid.org/0009-0001-7502-9233>
 Raju mollick: <https://orcid.org/0009-0002-7906-317x>
 Nishat Sultana: <https://orcid.org/0009-0001-2920-5216>

Article Info

Received on: 7 March 2025

Accepted on: 24 May 2025

Published on: 1 July 2025

References

- Chong Y, Park G, Cha HJ, Kim HJ, Kang CS, Abdul-Ghafar J, Lee SS. A stepwise approach to fine needle aspiration cytology of lymph nodes. *Journal of Pathology and Translational Medicine*. 2023 Jul 11;57(4):196-207.
- Zhou J, Li F, Meng L, et al. Fine needle aspiration cytology for lymph nodes: a three-year study. *Br J Biomed Sci* 2016; 73: 28–31.
- Cibas ES, Ducatman BS. *Cytology: diagnostic principles and clinical correlates*. 5th ed. Maryland Heights: Elsevier Inc, 2020.
- Rammeh S, Romdhane E, Sassi A, et al. Accuracy of fine-needle aspiration cytology of head and neck masses. *Diagn Cytopathol* 2019; 47: 394–9.
- Padia B, Dhokiya M. A study of FNAC of head and neck lesions at a tertiary care centre. *Trop J Path Micro*. 2018;4(8):592-6.
- Kaur A, Poonia S, Singh K, Kaur D, Madhukar M, Godara R. Cytomorphological spectrum of head and neck lesions by fine needle aspiration cytology in a tertiary care center. *Journal of Pharmacy and Bioallied Sciences*. 2023 Jul 1;15(Suppl 1):S315-7.
- Pathak R, Prasad KB, Rauniyar SK, Pudasaini S, Pande K, Koirala S, Kafle S, Jha A, Chalise S, Basnyat AS. Fine needle aspiration cytology of head and neck lesions and its correlation with histopathology. *Journal of Pathology of Nepal*. 2016 Sep 24;6(12):985-9.
- Deval N. Patel, Parth B. Patel, Himani V. Patel TJ, Gandhi. Fine needle aspiration cytology role in head and neck lesions. *Int Arch Inte grated Med*. 2015;2(8):99–104.
- Pawde Y, Kathale S. Fine Needle Aspiration Cytology As a Diagnostic Tool in Head and Neck Lesions. *J Evol Med Dent Sci J Evol Med Dent Sci*. 2014;3(45):11072–9.
- Bhagat VM, Tailor HJ, Saini PK, Dudhat RB, Makawana GR, Ravi M. Fine Needle Aspiration Cytology in NonThyroidal Head and Neck Masses-a Descriptive Study in Tertiary Care Hospital. *Natl J Med Res*. 2013;3(3):273–6.
- Gogoi G, Borgohain D. Fnac on Palpable Neck Masses – A Hospital Based Study. *Indian J Appllied Res*. 2016;6(4):39–42.
- Singal P, Ms B, Kharbanda J, Ps S. Efficacy of fine needle aspiration cytology in Head and Neck lesions. *IJMDS*. 2014;3(2):421–30.
- Shekhar H, Kaur A, Agrawal P, Pancharia A, Jadeja P. Fine needle aspiration cytology in head and neck swellings: a diagnostic and therapeutic procedure. *Int J Res Med Sci* 2014;2:1667-71.
- Sreedevi P, Kishore Kumar, Parankusa N C. Diagnostic Role of FNAC in evaluation of Head and Neck lesions. *Journal of Medical and Dental sciences*. 2016;15(9):11-13.
- Kaur A, Poonia S, Singh K, Kaur D, Madhukar M, Godara R. Cytomorphological spectrum of head and neck lesions by fine needle aspiration cytology in a tertiary care center. *Journal of Pharmacy and Bioallied Sciences*. 2023 Jul 1;15(Suppl 1):S315-7.
- Goswami RR, Baruah D, Devi G. FNAC spectrum of head and neck lesions-a retrospective study. *J Evid Based Med. Healthc*. 2016;3(13):400-5.

17. Deval N Patel, Parth B Patel, Himani V Patel. Fine needle aspiration cytology role in head and neck lesions. IAIM 2015;2(8):99-104.

18. Arvind Kumar B. Sangavi, Inder Raj Itagi, Suhas Y. Choudhari, Venkatesh U. Evaluation of FNAC of head and neck swellings: a retrospective study. Int J Otorhinolaryngol Head Neck Surg. 2018 Jan;4(1):189-192.

19. Md Hamed Altaf Mali, Arpitha K., Radha Angadi, Girish Ronad, Sanjeev Navadagi. Correlation Study of Cytomorphology, Histopathology and Immunohistochemistry in the Diagnosis of Lymphadenopathy. J Evid Based Med Healthc, pISSN - 2349-2562, eISSN - 2349-2570 / Vol. 8 / Issue 18 / May. 03, 2021.