

Case Report

Tradition of Adenoidectomy in children with recurrent URTI's

Karim R¹, Chowdhury AR², Hossain D³.

urgent (surgery) surgical interference doesn't bring any benefit over initial conservative treatment.

Although recurrent upper respiratory tract infection in childhood are self limiting with high rates of spontaneous recovery, they are associated with high health care costs and frequent visit to the doctor. Most of these infections are caused by viruses¹. Generally, they are associated with little morbidity but predispose children to complications such as otitis media, tonsillitis and sinusitis, which cause further morbidity. About a fifth of preschool children have recurrent upper respiratory tract infections, and many of these are referred to ear, nose and throat surgeons for upper airway surgery, including adenoidectomy.

Adenoidectomy restores normal breathing through nose if the hypertrophied adenoid obstructs the nasopharynx². It can also improve hearing in children with otitis media, but it has yet to show benefit on the occurrence of otitis media, physician visits, antibiotic prescription, or days with symptoms of upper respiratory infections in children under 2 years of age.

In the linked study, van den Aardweg and colleagues assessed the effectiveness of adenoidectomy in children selected for surgery on the basis of upper respiratory infections alone. The randomized controlled trial of 111 children aged 1-6 years compared immediate adenoidectomy, with or without myringotomy, with initial watchful waiting³. The primary outcome was the total number of recurrent upper respiratory tract infections defined as two or more of the following symptoms -fever, diary scored stuffiness or breathing through the mouth, nasal discharge, or cough during the two years of follow up. The study found no significant difference between the groups⁴. Furthermore, no difference between groups were found in the secondary

outcomes of total number of days with upper respiratory tract infections a year, days with fever, or health related quality of life.

The authors concluded that immediate surgery for upper respiratory tract infections has no benefit over watchful waiting currently, evidence on the benefits of adenoidectomy and internationally accepted guidelines are lacking in children with upper respiratory tract infection without additional symptoms of otitis media or sleep apnoea⁵. A symptomatic review in 2010 of adenoidectomy in children identified only two randomized controlled trials that included upper respiratory tract infections as an outcome. One was methodologically weak and the other included only children who primarily had otitis media.

The generalisability of the results warrants consideration⁶. Ear, nose and throat surgeons in the respective hospital diagnosed and selected eligible children. Because there are no internationally accepted guidelines for the diagnosis and treatment of upper respiratory tract infections, diagnosis is likely to vary between doctors⁷. Furthermore, rates of adenoidectomy varies considerably nationally and internationally and rates for recurrent upper respiratory infections alone have been estimated to be 12% in the United States and 60% in the Netherlands over the past decade.

Rates of upper respiratory tract surgery vary nationally and internationally⁸. Clinicians seem to be similar in their uncertainty regarding indications for procedure and variations in surgery rates among countries appear to be more characteristic for the procedure than for the country in which it is performed.

1. Dr. Rezaul Karim, MBBS, MS(ENT), Diploma in Aviation medicine (Pakistan), KYAMCH, Sirajgonj.

2. DR. Abu Reza Chowdhury Assistant Professor, East West Medical College, Dhaka.

3. Dr. Delowar Hossain, Associate Professor, BSMMU, Dhaka.

To date, no consensus exists to help surgeons decide which children with upper respiratory tract infection will benefit from adenoidectomy⁹. Accordingly, this lack of exact criteria for surgery may have affected the selection of children to surgery in the current study and also the primary and secondary outcomes in both study groups if the disease severity varied in the two treatment arms¹⁰. Further studies are needed to replicate the authors findings to refine the study design in terms of higher specificity in diagnostic inclusion criteria for surgery and finally to increase the sample size so that comparisons between different age groups can be made.

An important finding from van den Aardweg and colleagues study was the reduction in upper respiratory infections are self limiting and decrease with age¹¹. It also highlights the difficult in selecting children for surgery because symptom severity often reduces as the child grows older. Information and knowledge sharing are becoming increasingly important in childhood illness¹². Because a common genetic liability for upper air way infections is well known, and treatment for upper respiratory infections primarily aims to prevent complications such as tonsillitis, sinusitis, and otitis media, clinicians should be aware that children presenting with a family history of upper airway infection may be more vulnerable to recurrent infections careful follow up and a strategy if watchful waiting seem prudent in the light of this study, especially because the incidence of upper respiratory infections declines with age.

1. Nyquist AC, Gonzales R, Steiner JF, Sande MA, Antibiotic prescribing for children with colds, upper respiratory tract infections, and bronchitis. *JAMA* 1998;279:875-7.
2. Schilder AG, Zielhuis GA, Straaman HS, van den Broek p. An epidemiological approach to the etiology of middle ear disease in the Netherlands. *Eur Arch Otorhinolaryngol* 1992;249:370-3.
3. Haapkyla Karevold G, Kvaerner KJ, Pitkaranta A. Trends in otitis media sugery: a decrease in adenoidectomy. *IntJ Pediatr Otorhinolaryngol* 2008;72:1207-13.
4. Kvaerner KJ, Nafstad P, Jaakkola JJ. Otorhinolaryngol surgery and upper respiratory tract infections in children: an epidemiological study. *Ann otol Rhinolla Laryngol* 2002;111:1034-9.
5. Koiuvunen P, Uhari M, Luotonen J, Kristo A, Raski R, Pokka T et al. adenoidectomy versus chemoprophylaxis and placebo for recurrent acute otitis media in children aged under 2 years: randomized controlled trial. *BMJ* 2004;328:487.
6. Van den Aardweg MTA, Boonaker CWB, Rovers MM, Hoes AW, Schilder AGM, Effectiveness of adenoidectomy in children with recurrent upper respiratory tract infections: open randomized controlled trial. *BMJ* 2011;343:d5154.
7. Van den Aardweg Mt, Schilder AG, Herkert E, Booncker CW, Rovers MM. adenoidectomy for recurrent or chronic nasal symptoms in children. *Cochrane Database Syst Rev* 2010;1:CD008282.
8. McPhersin K, Wennberg JE, Hovind OB, Clifford P. Small area variations in the use of cammon surgical procedures: an international comparison of New England, England, and Norway. *N Eng J Med* 1982;307:1310-4.
9. Van den Akker EH, Hoes AW, Burton MJ, Schilder AR. Large international difference in (adeno) tonsillectomy rates. *Clin Otolaryngol Allied Sci* 2004;29:161-4
10. Bhattacharyya N, Lin HW. Changes and consistencies in the epidemiology of pediatric adenotonsillar surgery, 1996-2006. *Otolaryngol Head neck Surg* 2010;143:680-4.
11. Siegel RM. Acute otitis media guidelines, antibiotic use, and shared medical decision-making. *Pediatrics* 2010;125:384-6.
12. Kvestad E, Kvaerner KJ, Roysamb E, Tambs K, Harris JR, Magnus P. Recurrent otitis media and tonsillitis: Common disease predispositions. *Int J Pediatr Otorhinolaryngol* 2006;70:1561-8.