

**Chemical and Biological Investigation of
Leaves of *Polygonum plebejum***Md. Kamrul Hasan^{1*}, Abul Kalam Lutful Kabir², Sabyasachy Mistry¹Department of Natural Science, Stamford University Bangladesh¹
51 Siddeswari Road, Dhaka-1217Department of Pharmacy, Stamford University Bangladesh²***Corresponding Author**Md. Kamrul Hasan
Lecturer, Department of Natural Science
Stamford University Bangladesh
Contact No.: +8801552 304378
E-mail: kamrul_du79@yahoo.com

Received- 22 May, 2009

Accepted for Publication- 15 December, 2009

ABSTRACT

The cytotoxic and antimicrobial activity of methanol crude extract and column fractions of the extract of the leaves of *P. plebejum* were examined by brine shrimp lethality bioassay and disc diffusion method respectively. The extracts showed significant cytotoxic as well as antimicrobial activities. Silica gel column chromatography of methanol extract of *P. plebejum* afforded a steroid. The structure was elucidated on the basis of spectral analysis, including ¹H NMR and ¹³C NMR and also by comparing with data in the literature.

Key words: *Polygonum plebejum*, *Polygonaceae*, Stigmasterol, Cytotoxicity.

INTRODUCTION

The natural products, mainly of plant origin, have been used for treatment of diseases, and an impressive number of drugs have been isolated from natural sources. According to WHO about 4 billion people in the world rely on plant a sources of drugs (N.R. Fransworth, 1988). The plant kingdom represents reservoir of new chemical compounds (M. Hamvergur et al., 1991) such as glycosides, steroids, alkaloids, terpenoids, flavones, etc. are mainly responsible for their various therapeutic properties and pharmacological actions. Research studies leading to extraction, isolation and biological study of plant constituents have now formed the major field of study. The plant *Polygonum plebejum* belongs to the family *Polygonaceae*, locally known as Chemti Sag, Small Knotweed in English is a prostrate, densely branched, annual herb (Ghani A., 1998). The main constituents of the leaves of *P. plebejum* are flavonoids (13.03%), saponin (8.9%), steroids (4.8%) and alkaloids (0.7%) (Scalbcrt A. 1997). We investigated this plant and isolated a steroid. The purpose of this study is to identify and characterize the bioactive principles from the leaves of *P. plebejum*.

MATERIALS AND METHODS

Melting point of the compound was determined by using an electrothermal melting point apparatus (Mel-temp, OGAWA SEIKI CO., Japan). Shimadzu IR-470 spectrophotometer was used for recording IR (KBr pellets). The ¹H- and ¹³C- NMR spectra were recorded on 400 MHz (Bruker DPX400) NMR spectrometer using CDCl₃ (7.25/77.03) as the internal reference. TLC was carried out on pre-coated silica gel 60 PF₂₅₄ on aluminium sheets. For column chromatography silica gel, G-60 (70-230 mesh) was used.

Extraction and Isolation

The fresh leaves of *Polygonum plebejum* were collected from local market, Dhaka. The leaves were dried under 40°C and ground into coarse powder. The powdered sample (400 gm) was separately extracted with petroleum ether, dichloromethane and ethyl acetate in Soxhlet apparatus at 50°C. The extractive free powder was then extracted repeatedly with methanol in a reflux condition for half an hour. The methanol extract (M) was evaporated to dry mass (6.02 g) by using a rotary evaporator at 40°C and then subjected to column chromatography. The column was eluted with 100% pet-ether, dichloromethane and methanol in order of increasing polarity. The eluted sample was monitored by TLC, samples having similar TLC behavior was combined together to give twelve different fractions, C₁ to C₁₂ (15, 20, 13, 40, 18, 15, 10, 50, 90, 80, 60

and 900 mg, respectively). Among them the fraction C₆ (15 mg) was found to give single spot on TLC plate. So attempt was taken to characterize this compound.

Test for alcohol

Four gram of ceric ammonium nitrate was dissolved in 10 mL⁻¹ of 2N HNO₃ on mild heating. A few crystals of C₆ were dissolved in 0.5 mL⁻¹ of dioxane. The solution was added to 0.5 mL⁻¹ of ceric ammonium nitrate reagent, diluted to 1 mL⁻¹ with dioxane and shaken well. The solution was developed yellow to red color indicating the presence of an alcoholic hydroxyl group.

Test for steroid

Salkowski Reaction: A few crystals of C₆ were dissolved in chloroform and a few drops of concentrated sulfuric acid were added to the solution, a reddish color was seen in the upper chloroform layer.

Liebermann-Burchard reaction: A few crystals of C₆ were dissolved in chloroform and a few drops of concentrated sulfuric acid were added to it followed by the addition of 2-3 drops of acetic anhydride. The solution turned violet, blue and finally green.

Biological screening

Crude extract of methanol, two column fractions of methanol extract (C₃, C₉) and pure compound (C₆) were screened for their possible cytotoxicity by brim shrimp lethality bioassay and antimicrobial activity.

Cytotoxicity study

Brine shrimp lethality bioassay (Persoone, 1980) technique was applied for the determination of cytotoxic property of crude extract of methanol, two column fractions (C₃, C₉) and pure compound C₆ of the leaves of the plant *P. plebejum*.

Preparation of positive control group

Vincristine sulphate was used as the positive control. Measured amount of the vincristine sulphate was dissolved in DMSO to get an initial concentration of 20 µg/ml from which serial dilutions are made using DMSO to get 10 µg/ml, 5 µg/ml, 2.5 µg/ml, 1.25 µg/ml, 0.625 µg/ml, 0.3125 µg/ml, 0.15625 µg/ml, 0.078125 µg/ml and 0.0390 µg/ml. Then the positive control solutions are added to the pre-marked vials containing ten living brine shrimp nauplii in 5 ml simulated sea water to get the positive control groups.

Preparation of negative control group

30 µl of DMSO was added to each of three pre-marked glass vials containing 5 ml of simulated seawater and 10 shrimp nauplii to use as control groups. If the brine shrimps in these vials show a rapid mortality rate, then the test is considered as invalid as the nauplii died due to some reason other than the cytotoxicity of the compounds.

Counting of nauplii

After 24 hours, the vials were inspected using a magnifying glass and the number of survived nauplii in each vial were counted. From these data, the percent (%) of lethality of the brine shrimp nauplii was calculated for each concentration.

Antimicrobial assay

The disc diffusion method (Bauer et al., 1966) was used to test antimicrobial activity against thirteen bacteria and three fungi (Table-2). Solutions of known concentration (mg/ml) of the test samples were made by dissolving measured amount of the samples in calculated volume of solvents. Dried and sterilized filter paper disc (6 mm diameter) were then impregnated with known amounts of the test substances using micropipette. Discs containing the test material were placed on nutrient agar medium uniformly seeded with the test microorganisms. Standard antibiotic discs (kanamycin 30 µg/disc) and blank discs (impregnated with solvents) were used as a positive and negative control. These plates were then kept at low temperature (4°C) for 24 h to allow maximum diffusion. There is a gradual change in concentration in the media surrounding discs. The plates were then incubated at 37°C for 24 h to allow maximum growth of the organisms. The test materials having antimicrobial activity inhibited the growth of the microorganisms and a clear, distinct zone of inhibition was visualized surrounding the medium. The antimicrobial activity of the

test agent was determined by measuring the diameter of zone of inhibition expressed in millimeter. The experiment is carried out three times and the mean of the reading is required (Bauer et al., 1966).

RESULTS AND DISCUSSION

Compound C₆ was a white amorphous solid, soluble in dichloromethane and chloroform (Rowshanul MH et al., 2007), m.p. 155-160°C and showed positive results in alcohol test and in all the tests of steroids. IR (KBr) ν_{\max} : 3450, 2920, 2850, 1655, 1455, 1365 and 1040 cm^{-1} . The ¹H-NMR and ¹³C-NMR data C₆ are summarized in Table-1.

Cytotoxicity study

Following the procedure of Mayer (Larson, 1988), the lethality of M, C₃, C₉ and C₆ to brine shrimp was determined on *A. salina*. The LC₅₀ obtained from the best-fit line slope were found to be 2.82 $\mu\text{g/ml}$, 1.26 $\mu\text{g/ml}$, 3.98 $\mu\text{g/ml}$ and 1.20 $\mu\text{g/ml}$ for the test samples of M, C₃, C₉ and C₆ respectively.

Antimicrobial assay

The methanol crud extract (M), column fraction C₃, column fraction C₉ and pure compound C₆ of the leaves of *P. plebejum* were screened against thirteen bacteria and three fungi (Table 2) for antimicrobial activities by disc diffusion method.




Fig. 1: The chemical structure of C₆.

Table -1: ^1H -NMR and ^{13}C -NMR data of C_6

Position	δ_{H}	δ_{C}
1		32.9
2		34.5
3	3.25 (tdd j = 4.5 MHz)	79.0
4		42.0
5		154.6
6	5.14 (1H, m)	124.5
7		31.2
8		28.7
9		42.0
10		39.6
11		19.4
12		31.9
13		40.8
14		47.7
15		21.4
16		21.3
17		48.2
18	1.07 (3H, s)	18.3
19	1.26(3H, s)	18.2
20		33.4
21	0.91(3H, s)	17.4
22	4.62 (1H, m)	107.1
23	4.16 (1H, m)	139.5
24		47.7
25		30.6
26	1.01 (3H, s)	20.2
27	1.00(3H, s)	20.2
28		25.4
29	0.97 (3H, s)	12.2

δ_{H} = chemical shift values in ^1H -NMR spectrum; δ_{C} = chemical shift values in ^{13}C -NMR spectrum

Table 2: Antimicrobial activities of the different parts of methanol extract of the leaves of *Polygonum plebejum*.

Test Bacteria and Fungi	MCE (400 µg/disc)	CF-C ₃ (400 µg/disc)	CF-C ₉ (400 µg/disc)	PC (400 µg/disc)	Kanamycin (30 µg/disc)
Bacteria					
BC	10	10	9	10	35
BM	10	10	-	10	35
BS	9	10	-	10	36
SA	10	9	8	10	35
SL	10	10	8	9	35
EC	12	10	-	8	35
PA	10	8	-	9	36
SP	8	12	9	12	36
ST	10	10	10	10	36
VM	10	10	9	10	35
VP	8	9	-	8	35
SD	12	12	-	10	36
SB	10	12	10	12	35
Fungi					
CA	10	12	-	12	35
AN	12	12	12	12	35
SC	12	10	9	10	35

BC=*Bacillus cereus* (BTCC-19), BM=*Bacillus megaterium* (BTCC-18), BS=*Bacillus subtilis*, SA=*Staphylococcus aureus*, SL=*Sarcina lutea*, EC=*Escherichia coli* (BTCC-172), PA=*Pseudomonas aureus* (BTCC-1252), SP=*Salmonella paratyphi*, ST=*Salmonella typhi*, VM=*Vibrio mimicus*, VP=*Vibrio parahemolyticus*, SD=*Shigella dysenteriae*, SB=*Shigella boydii*, CA=*Candida albicans*, AN=*Aspergillus niger*, SC=*Sacharomyces cerevacaee*, MCE=Methanol Crud Extract, CF=Column Fraction, PC=Pure Compound, "-" indicates no sensitivity.

From the positive tests of for steroids and alcohols given by the C₆ was assumed to be a sterol. The melting point of C₆ (160°C) was in a good agreement with the melting point of stigmasterol in the literature (Holland et al., 1976). In ¹H-NMR spectrum of C₆, H-3 proton appeared as a triplet of a double of a double doublet (tdd) at δ 3.25 (J = 4.5 and 1.1 MHz) and H-6 olefinic proton showed a multiplet at δ 5.14. Two olefinic protons appeared doenfield at δ 4.14 (m) and δ 4.61 (m) which were identical with the chemical shift of H-22 and H-23, respectively of stigmasterol (Li et al., 2006). Six methyl protons also appeared at δ 1.07, δ 1.26, δ 0.91, δ 1.01, δ 1.00 and δ 0.97. ¹³C-NMR data of C₆ was also quite similar with the data in the literature of sigmasterol (Conolly and Hill, 1994)

Cytrocity study

The comparison with positive control (vincristine sulphate, LC₅₀ value 0.33 µg/mL), the pure compound as well as crude extract and column fractions showed potential cytotoxic activities.

Antimicrobial assay

Pure compound C₆ showed poor activity but crude extract and column fractions C₆ and C₉ exhibited moderate to significant activity against most of the test bacteria and fungi.

CONCLUSION

From these physical, chemical and spectral evidences C₆ was confirmed as stigmasterol (Fig. 1). The cytotoxicity exhibited by the pure compound and crude methanol extract of leaves is promising and they might have antitumor or pesticidal compounds (Meyer et al., 1982). However, this can not be confirmed without further higher and specific tests.

REFERENCES

- M. Hamvergur and K. Hostmann (1991) "Bioactivity in Plants: The Link Between Photochemistry and Medicine", *Phytochemistry*, 30(12) 3864-3874.
- Conolly, J.D. and R.A. Hill (1994) Dictionary of Natural Products, Chapman and Hall, pp: 4506.
- Ghani A. (1998) Medicinal Plants of Bangladesh, 1st Ed. pp. 63. The Asiatic Society of Bangladesh, Dhaka.
- Holland, H.L., P.R.P. Diakow and G.J.Taylor (1976). Microbial hydroxylation of steroids. *Can. J. chem.*, 56: 3121.
- N.R. Fransworth, *ibid.* (1988) pp. 83-97.
- Li, C., P.B. Bu, D.K. Yue and Y.F. Sun (2006) Chemical constituents from roots of *Ficus hirsuta*, *Zhongguo Zhong. Yao. Za. Zhi.*, 31: 131-133
- Scalbrt A. (1997) Medicinal plants, their bioactivity, screening and evaluation, proceeding of their international workshop. Lucknow (India) December 2-5, XL VII.
- Persone G. (1980) Proceeding of the International Symposium on Brine Shrimp, *Artemia salina*, Vols. 1-3. Universa Press, Witteren, Belgium.
- Bauer AW, Kirby WMM, Sherris JC, Turck M. (1966) Antibiotic susceptibility testing by a standardized single disc method. *Am. J. Clin. Pathol.* 45: 493-496
- Rowshanul MH, Farjana N, Matiar R, Ekramul MH, Rezaul MK. (2007) Isolation of Stigmasterol and β -Sitosterol from methanolic extract of root bark of *Calotropis gigantea* (Linn) *Pakistan Journal of Biological Sciences* 10 (22): 4174-4176
- Larson RA. (1988). The antioxidants of higher plants. *Phytochemistry* 27: 969-978
- Meyer BN, Ferringm NR, Puam JE, Lacobsen LB, Nichols DE, MeLaughlin JL. (1982) Brine shrimp: a convenient general bioassay for active constituents. *Planta Medica.* 45: 31-32.